

Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0101	197	93.4	0.0	2.0	0.0	0.0	4.6	0.0
0102	652	89.9	0.9	4.4	0.2	0.3	4.3	0.0
0103	657	90.7	0.2	3.2	0.0	0.0	5.9	0.0
0104	210	91.9	0.0	2.9	0.0	0.5	4.8	0.0
0250	252	78.2	0.0	0.4	0.4	0.0	20.6	0.4
0251	186	80.1	0.0	0.5	2.7	1.1	15.6	0.0
0252	222	45.0	0.0	2.7	0.9	0.9	50.5	0.0
0253	295	66.1	0.3	1.0	3.7	0.0	28.8	0.0
0401	821	92.2	0.2	1.9	0.0	0.0	5.6	0.0
0402	613	82.7	0.5	2.3	0.3	0.0	14.2	0.0
0403	228	95.2	0.0	2.6	0.0	0.4	1.8	0.0
0404	240	90.8	0.0	2.5	0.4	0.0	6.3	0.0
0501	913	91.6	0.4	3.0	0.2	0.1	4.7	0.0
0502	4	*	*	*	*	*	*	*
0503	292	88.4	0.0	6.5	0.0	0.3	4.8	0.0
0504	898	85.9	0.6	2.4	1.1	1.2	8.8	0.0
0505	616	85.4	0.3	4.2	0.2	0.0	9.9	0.0
0506	338	92.0	0.0	3.8	0.0	0.0	4.1	0.0
0650	1,590	82.6	3.7	4.2	1.1	0.9	7.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0651	1,041	87.9	0.1	5.3	0.1	0.3	6.1	0.2
0652	530	91.3	0.2	4.2	0.0	0.0	4.3	0.0
0653	766	96.3	0.3	2.1	0.0	0.3	1.0	0.0
0654	667	93.6	0.7	1.6	0.0	0.0	4.0	0.0
0655	592	92.9	0.2	1.2	0.2	0.5	5.1	0.0
0657	127	96.1	0.0	0.8	0.0	0.0	3.1	0.0
0701	674	89.5	0.9	2.5	0.0	3.6	3.6	0.0
0702	144	87.5	0.0	0.7	3.5	0.0	8.3	0.0
0703	357	88.5	0.0	5.0	1.1	0.0	5.0	0.3
0704	143	90.9	0.0	4.2	1.4	0.0	3.5	0.0
0705	284	93.0	0.0	1.1	0.0	0.0	6.0	0.0
0850	1,768	80.8	4.3	7.1	0.1	0.5	7.1	0.1
0851	1,463	91.3	0.5	4.5	0.0	0.2	3.2	0.3
0901	2,767	69.7	1.7	16.4	0.0	0.0	11.9	0.1
0902	143	83.9	0.0	6.3	0.7	0.0	9.1	0.0
1051	2,154	85.5	0.7	6.3	0.3	0.4	6.8	0.0
1101	2,899	30.8	57.2	5.1	0.2	1.0	5.5	0.1
1102	1,383	66.1	10.7	6.9	0.4	1.2	14.7	0.1
1106	174	77.6	1.7	13.2	0.0	0.0	6.3	1.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1250	1,755	78.8	0.6	16.7	0.0	0.5	3.4	0.1
1251	1,447	91.2	0.2	4.5	0.1	0.1	3.9	0.0
1253	2,441	69.2	3.1	21.1	0.0	0.1	6.6	0.0
1350	3,845	50.1	18.5	12.4	0.2	4.1	14.8	0.0
1351	218	86.7	1.4	1.8	0.0	0.9	9.2	0.0
1352	735	55.0	23.9	8.2	0.0	0.4	12.5	0.0
1401	357	82.4	0.6	8.4	0.6	0.0	7.8	0.3
1402	273	49.5	0.4	32.6	7.0	0.4	9.2	1.1
1403	58	*	*	*	*	*	*	*
1404	29	*	*	*	*	*	*	*
1501	1,884	39.8	7.7	46.0	0.1	0.0	6.2	0.2
1503	2,056	53.0	6.6	32.0	0.2	0.3	7.7	0.3
1504	3,932	39.1	4.0	48.4	0.0	3.0	5.5	0.0
1505	1,477	83.3	2.1	9.8	0.0	0.6	4.1	0.0
1506	1,087	74.8	1.1	18.6	0.2	0.0	5.2	0.1
1507	1,040	85.7	1.2	5.9	0.1	0.7	6.5	0.0
1508	230	0.0	0.0	98.3	0.0	0.0	0.0	1.7
1509	1,057	50.9	2.6	44.6	0.0	0.1	1.9	0.0
1511	1,155	88.8	1.4	6.0	0.1	0.3	3.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1512	666	84.1	0.3	8.7	0.0	0.0	6.9	0.0
1701	1,171	45.4	0.4	3.7	17.3	0.9	32.2	0.0
1950	1,019	89.4	0.4	6.2	0.0	0.1	3.9	0.0
1952	247	89.1	0.0	5.7	0.4	0.4	3.2	1.2
1953	510	85.9	0.8	9.0	0.2	0.2	3.9	0.0
1954	957	81.4	0.0	16.0	0.0	0.0	2.6	0.0
1955	266	79.3	0.4	17.3	0.0	0.4	2.6	0.0
1957	987	82.1	0.8	12.5	0.1	0.1	4.5	0.0
1959	80	*	*	*	*	*	*	*
2150	1,134	82.5	0.0	2.7	1.1	0.1	13.6	0.1
2151	414	88.9	0.2	3.6	0.5	0.0	6.8	0.0
2251	678	85.0	0.0	4.7	0.0	0.4	9.3	0.6
2252	358	91.9	0.0	2.5	0.3	0.0	5.3	0.0
2551	253	26.1	65.2	2.0	0.4	0.0	6.3	0.0
2552	9	*	*	*	*	*	*	*
2554	234	79.5	10.7	3.8	0.0	0.4	5.6	0.0
2555	10,731	47.0	39.0	5.6	0.1	0.3	8.1	0.0
3001	1,636	11.9	66.4	17.7	0.0	0.4	3.7	0.0
3002	460	17.4	47.8	29.6	0.0	0.2	5.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
3003	4	*	*	*	*	*	*	*
3004	283	0.0	0.0	100.0	0.0	0.0	0.0	0.0
3007	56	*	*	*	*	*	*	*
3009	17	*	*	*	*	*	*	*
3201	922	90.5	0.2	5.2	0.1	0.3	3.7	0.0
3301	5,454	31.1	28.3	23.1	0.0	4.4	13.0	0.0
3302	461	20.0	37.5	25.4	0.0	3.7	13.4	0.0
3304	276	39.9	19.6	11.2	0.0	24.3	5.1	0.0
3305	283	57.6	13.8	12.7	0.4	3.5	12.0	0.0
3306	461	87.2	1.5	4.6	0.0	0.2	6.3	0.2
3308	115	86.1	0.0	7.0	0.0	0.0	7.0	0.0
3309	96	*	*	*	*	*	*	*
3550	149	0.0	0.0	2.0	91.3	0.0	6.7	0.0
3551	44	*	*	*	*	*	*	*
3552	88	*	*	*	*	*	*	*
3701	1,847	82.8	0.3	13.4	0.1	0.1	3.4	0.0
3805	5,051	63.8	12.5	8.9	0.1	0.3	14.4	0.0
3950	2,997	57.1	17.3	10.8	0.0	0.8	13.9	0.1
3952	2	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
3954	104	72.1	2.9	6.7	0.0	6.7	11.5	0.0
4001	1,982	21.1	57.3	9.9	0.1	0.6	10.9	0.1
4002	1,855	39.0	23.7	20.4	0.1	0.6	16.2	0.0
4103	227	5.7	73.1	6.2	0.0	5.7	9.3	0.0
4106	3,427	11.3	37.7	43.3	0.0	0.2	7.5	0.0
4107	5,276	42.1	22.0	21.3	0.1	1.3	13.2	0.0
4109	1,960	81.7	1.3	8.9	0.0	0.2	8.0	0.0
4110	923	21.6	10.9	56.3	0.0	0.4	10.7	0.0
4112	1,070	0.1	0.3	99.5	0.0	0.1	0.0	0.0
4113	1,182	4.5	16.0	76.6	0.0	0.3	2.7	0.0
4120	3,444	43.9	8.0	35.4	0.2	2.1	10.4	0.0
4160	2,944	32.7	30.3	17.2	0.1	8.9	10.7	0.0
4351	979	85.2	0.1	8.5	0.0	0.1	6.1	0.0
4352	310	62.6	12.9	5.5	0.0	0.0	18.7	0.3
4353	806	64.5	0.4	31.1	0.0	0.1	3.8	0.0
4354	1,226	84.9	3.8	3.2	0.2	0.1	7.6	0.2
4355	509	73.3	0.2	12.4	0.0	0.0	14.1	0.0
4356	916	77.6	0.4	14.8	0.1	0.3	6.7	0.0
4358	336	3.6	0.0	95.8	0.0	0.0	0.3	0.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 7
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
4359	976	64.2	0.0	33.8	0.1	0.0	1.7	0.1
4360	309	86.4	0.0	11.3	0.0	0.0	2.3	0.0
4361	363	91.5	0.0	1.1	0.0	0.0	7.4	0.0
4362	525	90.3	0.0	4.4	0.0	0.2	5.1	0.0
4363	1,434	91.6	0.1	3.8	0.1	0.2	4.3	0.0
4401	2,027	84.4	0.2	7.2	0.0	0.0	8.2	0.0
4501	1,654	88.0	1.0	7.3	0.0	0.1	3.7	0.0
4502	3	*	*	*	*	*	*	*
4591	1	*	*	*	*	*	*	*
4601	2,292	70.3	1.0	21.6	0.0	0.0	6.9	0.1
4701	2,036	89.6	0.6	5.0	0.2	0.9	3.7	0.0
5050	2,457	57.9	26.6	3.1	0.1	1.8	10.4	0.0
5051	3,952	54.2	27.3	5.8	0.1	1.6	10.9	0.1
5052	3,896	56.5	21.6	2.8	0.0	9.7	9.3	0.0
5101	3,303	66.2	8.9	15.0	0.0	1.0	9.0	0.0
5202	969	83.7	0.7	3.2	0.6	0.0	11.8	0.0
5203	244	83.2	1.2	5.7	0.0	0.0	9.8	0.0
5204	186	76.9	1.1	4.8	0.0	0.0	17.2	0.0
5601	2,032	88.5	0.9	3.2	0.1	0.5	6.6	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 8
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
5801	2,813	78.3	4.9	8.2	0.0	0.1	8.3	0.1
5803	429	87.9	0.2	5.6	0.0	0.9	5.4	0.0
5804	167	85.0	1.8	8.4	0.0	0.0	4.8	0.0
6101	129	86.0	0.0	10.1	0.0	0.0	3.9	0.0
6102	400	80.5	1.8	7.8	0.3	0.0	9.8	0.0
6103	5,771	51.6	23.3	12.9	0.1	0.2	11.9	0.0
6104	691	24.7	52.7	12.2	0.1	0.1	10.1	0.0
6105	131	80.9	1.5	15.3	0.0	0.0	2.3	0.0
6106	4	*	*	*	*	*	*	*
6301	6,845	39.4	26.3	24.5	0.1	2.3	7.4	0.0
6312	4,207	39.2	49.3	3.2	0.1	1.7	6.5	0.0
6316	2,339	68.6	15.0	8.9	0.1	2.1	5.3	0.0
6320	385	63.4	27.5	2.9	0.0	2.6	3.6	0.0
6325	255	92.9	0.8	3.5	0.0	0.4	2.4	0.0
6330	639	22.7	53.7	16.7	0.0	0.0	6.9	0.0
6335	984	80.6	8.4	3.4	0.1	3.3	4.3	0.0
6340	213	37.6	21.1	33.8	0.0	3.8	3.8	0.0
6345	45	*	*	*	*	*	*	*
6350	429	54.5	34.0	2.8	0.0	0.7	7.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 9
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
7304	359	86.1	0.6	10.0	0.0	0.0	3.3	0.0
7306	4,011	40.3	32.0	19.6	0.0	0.4	7.6	0.1
7308	282	86.9	1.1	5.0	0.4	0.0	6.4	0.4
7310	76	*	*	*	*	*	*	*
7401	3,979	75.0	4.0	8.5	0.1	0.2	12.2	0.0
7601	1,099	87.3	0.1	9.3	0.0	0.2	3.2	0.0
7850	1,732	86.4	0.2	7.0	0.1	0.1	6.1	0.2
7852	162	92.6	0.0	1.9	0.0	0.0	5.6	0.0
8101	764	8.8	83.4	2.4	0.0	1.2	4.3	0.0
8102	136	3.7	2.9	93.4	0.0	0.0	0.0	0.0
8105	218	57.3	33.9	2.8	0.0	0.5	5.5	0.0
8106	179	19.6	60.3	7.3	0.0	0.6	12.3	0.0
8150	90	*	*	*	*	*	*	*
8151	4,361	32.3	39.1	12.9	0.2	2.2	13.2	0.2
8152	403	29.3	52.9	4.5	0.0	0.0	13.4	0.0
8153	49	*	*	*	*	*	*	*
8154	535	32.9	21.3	13.5	0.0	28.0	4.1	0.2
8155	136	49.3	28.7	7.4	1.5	5.9	7.4	0.0
8156	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 10
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
8157	33	*	*	*	*	*	*	*
8201	3,334	53.9	25.5	7.0	0.0	2.3	11.2	0.0
8202	5,291	85.5	7.7	4.1	0.1	0.8	1.8	0.0
8203	551	62.8	25.6	4.9	0.0	0.4	6.4	0.0
8250	1,354	26.8	57.9	5.4	0.1	1.5	8.3	0.0
8252	135	0.7	93.3	0.0	0.0	3.7	2.2	0.0
8255	3,873	56.1	24.4	10.0	0.1	1.8	7.6	0.0
8257	782	49.4	18.3	0.5	0.0	29.4	2.4	0.0
8258	1,796	62.8	22.9	6.0	0.1	0.6	7.6	0.1
8260	1,229	47.4	26.9	2.0	0.1	20.2	3.5	0.0
8901	1,795	0.8	94.9	0.7	0.1	0.8	2.7	0.0
8902	1,257	3.1	93.0	1.0	0.0	0.3	2.6	0.0
8903	2,081	1.3	95.4	0.9	0.0	0.5	1.9	0.0
8904	1,857	3.1	85.3	0.8	0.2	6.6	4.0	0.1
8905	4,862	42.3	26.0	25.4	0.0	1.8	4.5	0.0
9004	1	*	*	*	*	*	*	*
9010	6	*	*	*	*	*	*	*
9012	91	*	*	*	*	*	*	*
9013	170	0.0	95.9	2.9	0.0	1.2	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 11
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
9015	57	*	*	*	*	*	*	*
9016	291	0.0	96.6	0.0	0.0	0.3	3.1	0.0
9017	120	2.5	94.2	0.8	0.0	0.0	2.5	0.0
9018	196	1.0	91.3	2.6	0.0	0.0	5.1	0.0
9019	71	*	*	*	*	*	*	*
9021	160	0.0	95.6	1.3	0.0	0.0	3.1	0.0
9040	1	*	*	*	*	*	*	*
9041	103	0.0	97.1	2.9	0.0	0.0	0.0	0.0
9045	2,603	5.8	33.9	58.1	0.0	0.2	2.0	0.0
9050	2,432	12.4	48.8	37.1	0.0	0.2	1.4	0.0
9051	2,666	5.7	9.2	83.4	0.0	0.0	1.7	0.0
9053	3,092	0.9	95.3	1.2	0.0	0.8	1.7	0.0
9055	13	*	*	*	*	*	*	*
9058	1,346	63.6	28.9	0.7	0.1	5.5	1.2	0.0
9066	2,704	1.0	96.7	0.3	0.0	0.2	1.7	0.0
9068	1,913	1.4	95.7	0.7	0.1	0.4	1.8	0.0
9069	2,593	1.2	95.0	0.7	0.0	0.1	3.0	0.0
9070	2,400	2.0	84.9	0.3	0.0	9.5	3.3	0.0
9072	2,023	2.0	94.5	0.1	0.0	1.5	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 12
Run Date: 3 / 12 / 12

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
9074	1,547	0.6	95.7	0.4	0.1	0.2	3.0	0.0
9080	4,142	24.8	69.6	2.6	0.1	0.1	2.7	0.1
Unknown Clinics	534	33.3	21.2	20.2	6.7	2.8	15.7	0.0
Michigan	257,910	51.1	25.7	14.0	0.3	1.5	7.4	0.0
Nation (Prior Year)	8,672,965	32.5	19.1	39.7	0.9	2.5	2.7	2.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 13
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0101	197	28.4	19.8	51.8
0102	652	35.6	21.0	43.4
0103	657	33.5	19.9	46.6
0104	210	32.9	22.9	44.3
0250	252	27.0	23.0	50.0
0251	186	28.5	20.4	51.1
0252	222	17.6	30.6	51.8
0253	295	30.5	18.3	51.2
0401	821	31.5	21.8	46.7
0402	613	28.4	23.7	48.0
0403	228	25.0	23.2	51.8
0404	240	28.8	21.7	49.6
0501	913	31.3	21.1	47.5
0502	4	*	*	*
0503	292	32.2	21.9	45.9
0504	898	32.3	23.1	44.7
0505	616	35.6	16.9	47.6
0506	338	24.6	19.8	55.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 14
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0650	1,590	30.3	22.9	46.9
0651	1,041	30.3	21.9	47.8
0652	530	28.5	20.8	50.8
0653	766	25.7	22.7	51.6
0654	667	25.6	22.2	52.2
0655	592	33.8	20.4	45.8
0657	127	26.8	16.5	56.7
0701	674	31.9	20.6	47.5
0702	144	32.6	20.8	46.5
0703	357	34.2	18.5	47.3
0704	143	25.2	18.2	56.6
0705	284	32.7	16.9	50.4
0850	1,768	33.7	20.8	45.5
0851	1,463	29.6	20.4	50.0
0901	2,767	31.3	22.2	46.6
0902	143	28.7	28.0	43.4
1051	2,154	34.0	20.6	45.4
1101	2,899	31.8	22.0	46.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 15
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1102	1,383	32.5	23.6	43.8
1106	174	35.1	24.7	40.2
1250	1,755	31.9	18.3	49.8
1251	1,447	34.7	18.1	47.2
1253	2,441	32.8	19.7	47.5
1350	3,845	33.0	19.9	47.1
1351	218	28.4	21.6	50.0
1352	735	29.4	21.2	49.4
1401	357	27.5	21.0	51.5
1402	273	27.1	21.2	51.6
1403	58	*	*	*
1404	29	*	*	*
1501	1,884	27.2	21.1	51.8
1503	2,056	27.8	21.5	50.6
1504	3,932	27.1	24.6	48.3
1505	1,477	28.4	21.1	50.4
1506	1,087	29.0	21.2	49.9
1507	1,040	29.2	21.5	49.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 16
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1508	230	20.9	27.4	51.7
1509	1,057	23.8	22.5	53.6
1511	1,155	36.4	18.4	45.3
1512	666	28.2	21.3	50.5
1701	1,171	34.4	19.0	46.5
1950	1,019	37.5	19.2	43.3
1952	247	28.7	20.6	50.6
1953	510	29.6	20.2	50.2
1954	957	33.0	20.1	46.9
1955	266	31.2	17.3	51.5
1957	987	34.9	19.6	45.6
1959	80	*	*	*
2150	1,134	30.2	20.7	49.0
2151	414	30.7	20.5	48.8
2251	678	32.2	21.4	46.5
2252	358	29.3	16.8	53.9
2551	253	39.5	20.9	39.5
2552	9	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 17
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
2554	234	27.4	19.7	53.0
2555	10,731	40.4	18.4	41.2
3001	1,636	30.0	17.9	52.1
3002	460	28.3	18.7	53.0
3003	4	*	*	*
3004	283	24.7	13.8	61.5
3007	56	*	*	*
3009	17	*	*	*
3201	922	34.3	20.4	45.3
3301	5,454	36.4	20.2	43.4
3302	461	26.0	23.2	50.8
3304	276	29.0	18.8	52.2
3305	283	35.0	16.6	48.4
3306	461	31.0	21.0	47.9
3308	115	24.3	27.0	48.7
3309	96	*	*	*
3550	149	30.9	12.8	56.4
3551	44	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 18
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3552	88	*	*	*
3701	1,847	31.1	22.5	46.5
3805	5,051	32.4	21.1	46.5
3950	2,997	29.4	22.1	48.5
3952	2	*	*	*
3954	104	22.1	16.3	61.5
4001	1,982	33.6	19.1	47.3
4002	1,855	32.0	20.3	47.7
4103	227	22.5	18.9	58.6
4106	3,427	31.5	20.2	48.3
4107	5,276	33.6	19.9	46.6
4109	1,960	29.8	18.6	51.5
4110	923	23.8	22.3	53.8
4112	1,070	26.4	18.6	55.0
4113	1,182	31.1	18.4	50.5
4120	3,444	33.0	19.5	47.5
4160	2,944	34.3	20.6	45.1
4351	979	28.4	22.9	48.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 19
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4352	310	26.1	18.7	55.2
4353	806	25.2	24.1	50.7
4354	1,226	33.6	21.1	45.3
4355	509	25.1	24.4	50.5
4356	916	27.4	22.4	50.2
4358	336	19.6	24.1	56.3
4359	976	26.4	20.4	53.2
4360	309	32.0	16.2	51.8
4361	363	25.9	22.3	51.8
4362	525	26.1	22.9	51.0
4363	1,434	33.6	19.4	47.0
4401	2,027	34.0	21.3	44.6
4501	1,654	30.7	20.0	49.3
4502	3	*	*	*
4591	1	*	*	*
4601	2,292	36.3	19.9	43.8
4701	2,036	30.2	26.1	43.7
5050	2,457	35.1	22.7	42.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 20
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
5051	3,952	35.8	21.6	42.6
5052	3,896	35.2	22.8	42.1
5101	3,303	34.5	22.0	43.4
5202	969	34.1	21.9	44.1
5203	244	23.0	25.0	52.0
5204	186	28.0	20.4	51.6
5601	2,032	33.8	19.7	46.6
5801	2,813	34.6	21.4	43.9
5803	429	26.8	25.6	47.6
5804	167	22.8	21.0	56.3
6101	129	26.4	24.0	49.6
6102	400	24.3	18.8	57.0
6103	5,771	31.0	20.9	48.1
6104	691	47.5	16.4	36.2
6105	131	25.2	24.4	50.4
6106	4	*	*	*
6301	6,845	40.2	19.0	40.8
6312	4,207	41.0	18.6	40.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 21
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
6316	2,339	40.5	21.5	37.9
6320	385	27.5	17.1	55.3
6325	255	29.4	21.2	49.4
6330	639	33.3	17.7	49.0
6335	984	28.4	20.7	50.9
6340	213	32.4	13.6	54.0
6345	45	*	*	*
6350	429	29.6	24.0	46.4
7304	359	27.9	20.6	51.5
7306	4,011	34.3	20.2	45.5
7308	282	33.0	24.5	42.6
7310	76	*	*	*
7401	3,979	33.0	21.2	45.8
7601	1,099	33.8	20.0	46.2
7850	1,732	29.6	22.1	48.3
7852	162	32.1	19.8	48.1
8101	764	59.7	11.1	29.2
8102	136	25.7	13.2	61.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 22
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8105	218	51.8	15.6	32.6
8106	179	37.4	12.3	50.3
8150	90	*	*	*
8151	4,361	35.2	21.9	42.9
8152	403	49.9	22.8	27.3
8153	49	*	*	*
8154	535	34.0	16.6	49.3
8155	136	25.0	20.6	54.4
8156	31	*	*	*
8157	33	*	*	*
8201	3,334	43.2	23.9	32.9
8202	5,291	33.1	17.9	49.0
8203	551	37.9	23.4	38.7
8250	1,354	43.0	23.5	33.5
8252	135	40.7	11.9	47.4
8255	3,873	44.0	21.7	34.3
8257	782	30.2	21.6	48.2
8258	1,796	35.0	21.1	43.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 23
Run Date: 3 / 12 / 12

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8260	1,229	34.4	20.6	45.0
8901	1,795	37.9	18.9	43.2
8902	1,257	40.0	18.1	41.9
8903	2,081	37.0	18.3	44.7
8904	1,857	40.4	18.5	41.1
8905	4,862	32.3	18.4	49.3
9004	1	*	*	*
9010	6	*	*	*
9012	91	*	*	*
9013	170	41.2	19.4	39.4
9015	57	*	*	*
9016	291	39.5	17.5	43.0
9017	120	34.2	17.5	48.3
9018	196	37.8	19.9	42.3
9019	71	*	*	*
9021	160	36.3	16.9	46.9
9040	1	*	*	*
9041	103	18.4	7.8	73.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9045	2,603	30.7	21.4	47.8
9050	2,432	29.8	23.0	47.2
9051	2,666	30.3	18.0	51.7
9053	3,092	43.2	17.9	39.0
9055	13	*	*	*
9058	1,346	27.3	21.7	51.0
9066	2,704	36.2	20.5	43.3
9068	1,913	35.1	21.3	43.5
9069	2,593	43.2	21.0	35.8
9070	2,400	34.1	22.9	43.0
9072	2,023	37.8	21.7	40.5
9074	1,547	37.7	20.0	42.3
9080	4,142	32.6	22.1	45.4
Unknown Clinics	534	31.3	17.6	51.1
Michigan	257,910	33.8	20.6	45.6
Nation (Prior Year)	8,672,965	33.7	22.1	44.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 25
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0101	37	* *	* *	197	6.1(127)	2.0 (63)	7.6 (13)	102	16.7 (78)	9.8 (15)	139	14.4 (113)
0102	171	5.8 (10)	12.3 (112)	646	6.7(150)	4.3(157)	13.2(150)	285	17.2 (92)	16.5(135)	405	11.1 (81)
0103	165	6.7 (20)	6.7 (46)	650	6.5(143)	1.8 (48)	12.3(133)	301	21.9(151)	17.6(146)	430	7.9 (52)
0104	46	* *	* *	210	4.3 (39)	1.9 (59)	10.0 (58)	92	* *	* *	124	5.6 (29)
0250	49	* *	* *	249	4.0 (29)	3.2(132)	10.8 (87)	119	18.5(119)	16.8(139)	182	8.8 (59)
0251	30	* *	* *	185	3.2 (10)	0.0 (1)	15.7(176)	99	* *	* *	127	3.9 (12)
0252	33	* *	* *	220	5.0 (72)	1.4 (21)	14.1(165)	111	18.0(109)	19.8(152)	142	7.7 (50)
0253	58	* *	* *	293	2.4 (3)	9.6(183)	4.8 (1)	147	7.5 (2)	6.1 (3)	195	7.2 (45)
0401	177	5.6 (8)	13.0 (114)	816	5.3 (85)	1.8 (48)	11.4(107)	373	14.7 (38)	15.0(112)	541	3.9 (12)
0402	130	6.9 (23)	6.2 (34)	609	4.3 (39)	1.8 (48)	11.3(101)	299	16.4 (69)	14.7(108)	409	13.7 (107)
0403	41	* *	* *	223	3.6 (20)	1.8 (48)	14.3(169)	114	14.0 (32)	21.1(156)	152	0.7 (1)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 26
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
0404	43	* *	* *	237	3.0 (4)	1.7 (38)	11.0 (92)	117	17.1 (88)	15.4 (121)	152 10.5 (74)
0501	201	12.4 (105)	7.5 (69)	899	4.4 (46)	2.7 (109)	9.6 (49)	433	17.1 (88)	12.9 (66)	594 7.2 (45)
0502	0	* *	* *	3	* *	* *	* *	1	* *	* *	4 * *
0503	59	* *	* *	288	4.5 (47)	7.3 (178)	6.9 (8)	133	9.8 (5)	7.5 (7)	192 11.5 (85)
0504	180	8.9 (69)	9.4 (95)	889	4.8 (57)	1.8 (48)	11.2 (95)	413	18.9 (125)	12.3 (51)	604 7.3 (47)
0505	138	5.1 (5)	10.9 (108)	598	5.2 (79)	1.8 (48)	8.9 (32)	289	18.3 (117)	12.8 (64)	413 4.8 (19)
0506	62	* *	* *	334	3.9 (26)	5.1 (168)	8.7 (27)	188	14.9 (41)	12.2 (47)	238 5.0 (22)
0650	363	6.9 (23)	5.5 (27)	1,572	5.2 (79)	3.0 (125)	10.2 (67)	734	17.0 (86)	14.2 (95)	1,081 6.2 (34)
0651	230	6.1 (12)	7.0 (53)	1,029	5.5 (98)	2.0 (63)	11.3 (101)	489	14.9 (41)	15.3 (120)	676 15.8 (118)
0652	104	6.7 (20)	11.5 (111)	522	7.5 (163)	1.5 (27)	11.9 (118)	266	16.5 (73)	14.3 (97)	339 10.3 (70)
0653	138	8.7 (61)	8.7 (86)	760	5.1 (74)	2.2 (78)	12.5 (137)	386	18.1 (113)	16.3 (130)	525 11.0 (80)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 27
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
0654	136	8.1 (40)	7.4 (66)	658	4.1 (32)	2.0 (63)	10.5 (76)	340	17.9(108)	14.7(108)	453 16.8 (125)
0655	148	8.1 (40)	6.8 (48)	581	7.6(165)	2.1 (73)	9.5 (47)	265	15.1 (46)	14.3 (97)	371 3.5 (9)
0657	24	* *	* *	125	4.8 (57)	1.6 (34)	10.4 (73)	67	* *	* *	89 * *
0701	137	5.1 (5)	7.3 (63)	673	5.2 (79)	1.5 (27)	7.9 (15)	323	10.5 (6)	10.8 (21)	407 10.3 (70)
0702	29	* *	* *	144	7.6(165)	2.8(114)	9.7 (53)	68	* *	* *	88 * *
0703	74	* *	* *	354	4.2 (34)	2.3 (83)	11.6(112)	170	21.2(144)	17.1(140)	225 1.3 (2)
0704	28	* *	* *	142	4.2 (34)	2.1 (73)	9.9 (55)	84	* *	* *	85 * *
0705	53	* *	* *	282	6.7(150)	1.8 (48)	6.4 (5)	145	20.7(141)	4.1 (1)	163 2.5 (4)
0850	401	7.7 (35)	9.2 (93)	1,749	6.2(129)	2.6(100)	12.2(131)	789	16.9 (84)	16.3(130)	1,147 5.2 (23)
0851	283	7.4 (31)	8.1 (78)	1,425	4.1 (32)	2.7(109)	10.2 (67)	716	14.7 (38)	13.1 (72)	1,017 5.3 (25)
0901	602	9.1 (70)	9.0 (90)	2,747	6.2(129)	1.4 (21)	10.3 (71)	1,280	16.3 (66)	14.5(103)	1,882 9.1 (62)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 28
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
0902	24	* *	* *	141	9.9(181)	4.3(157)	12.1(128)	62	* *	* *	103 18.4 (131)
1051	556	10.3 (89)	10.8 (107)	2,085	4.8 (57)	6.9(177)	11.0 (92)	958	16.3 (66)	12.5 (58)	1,286 12.5 (92)
1101	710	11.5 (101)	7.2 (57)	2,848	6.5(143)	2.9(119)	10.4 (73)	1,329	15.6 (52)	14.1 (89)	1,928 10.4 (73)
1102	333	7.5 (33)	9.3 (94)	1,358	4.7 (56)	1.3 (16)	11.5(109)	590	16.6 (76)	15.9(126)	895 6.4 (37)
1106	41	* *	* *	171	3.5 (17)	0.0 (1)	8.2 (18)	73	* *	* *	104 4.8 (19)
1250	344	8.1 (40)	10.2 (103)	1,732	7.0(159)	2.0 (63)	14.3(169)	878	17.7(105)	14.6(105)	1,189 4.0 (14)
1251	308	8.8 (65)	8.4 (82)	1,432	5.4 (90)	1.5 (27)	8.9 (32)	676	17.2 (92)	11.5 (31)	966 2.3 (3)
1253	542	8.1 (40)	9.4 (95)	2,414	7.6(165)	2.6(100)	10.6 (79)	1,161	18.5(119)	13.0 (71)	1,671 11.5 (85)
1350	837	8.1 (40)	8.5 (83)	3,803	6.6(146)	2.5 (94)	14.3(169)	1,793	18.9(125)	16.6(137)	2,586 16.2 (122)
1351	45	* *	* *	216	4.6 (52)	1.9 (59)	10.6 (79)	110	12.7 (19)	13.6 (82)	154 11.7 (88)
1352	160	8.8 (65)	8.8 (89)	720	6.3(139)	2.5 (94)	11.3(101)	355	15.2 (48)	15.2(117)	490 19.2 (136)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 29
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
1401	66	* *	* *	353	4.8 (57)	4.5(163)	13.0(147)	183	16.4 (69)	15.8(125)	251 4.0 (14)
1402	48	* *	* *	266	3.4 (14)	1.9 (59)	13.9(161)	143	21.0(143)	12.6 (60)	202 13.4 (106)
1403	11	* *	* *	56	* *	* *	* *	28	* *	* *	42 * *
1404	4	* *	* *	28	* *	* *	* *	17	* *	* *	24 * *
1501	346	6.6 (17)	7.8 (73)	1,854	3.8 (23)	2.4 (87)	14.1(165)	966	17.1 (88)	15.9(126)	1,366 19.1 (135)
1503	389	8.2 (48)	7.2 (57)	2,028	4.2 (34)	2.5 (94)	11.3(101)	1,018	18.8(123)	13.3 (75)	1,401 30.5 (156)
1504	843	6.0 (11)	6.9 (50)	3,894	4.3 (39)	3.2(132)	11.3(101)	1,889	16.5 (73)	13.5 (79)	2,603 21.1 (143)
1505	299	8.4 (53)	11.4 (110)	1,459	6.2(129)	2.7(109)	12.1(128)	748	15.0 (44)	11.5 (31)	966 22.6 (147)
1506	224	7.1 (27)	7.1 (55)	1,076	5.7(106)	3.3(137)	10.5 (76)	537	19.9(133)	12.1 (43)	758 20.7 (140)
1507	215	9.3 (74)	10.7 (105)	1,022	3.8 (23)	3.3(137)	8.9 (32)	504	18.5(119)	10.1 (16)	689 30.8 (158)
1508	35	* *	* *	217	1.4 (1)	0.9 (10)	16.1(177)	111	30.6(159)	17.1(140)	145 20.0 (138)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 30
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1509	179	8.4 (53)	6.7 (46)	1,045	4.3 (39)	2.7(109)	13.7(157)	562	16.7 (78)	17.4(145)	758	22.0 (144)
1511	297	7.4 (31)	10.1 (101)	1,142	3.7 (21)	2.5 (94)	12.1(128)	525	16.6 (76)	15.4(121)	729	20.0 (138)
1512	125	8.8 (65)	7.2 (57)	660	3.3 (12)	2.9(119)	12.6(140)	341	19.1(127)	14.1 (89)	452	36.9 (163)
1701	260	6.5 (15)	14.2 (116)	1,159	4.5 (47)	3.0(125)	13.1(148)	536	17.5(102)	14.7(108)	750	5.3 (25)
1950	241	7.9 (36)	7.5 (69)	1,007	8.1(178)	1.5 (27)	11.5(109)	435	18.2(115)	15.9(126)	678	7.4 (49)
1952	45	* *	* *	243	4.5 (47)	1.6 (34)	9.9 (55)	121	13.2 (27)	13.2 (74)	169	5.3 (25)
1953	88	* *	* *	505	5.7(106)	2.4 (87)	11.5(109)	257	12.5 (15)	14.4(100)	347	5.2 (23)
1954	210	7.1 (27)	4.8 (14)	944	6.0(123)	2.6(100)	10.6 (79)	440	12.0 (11)	13.4 (77)	608	5.9 (31)
1955	52	* *	* *	265	4.9 (67)	2.3 (83)	12.5(137)	134	13.4 (28)	14.2 (95)	165	4.8 (19)
1957	220	8.2 (48)	6.8 (48)	976	5.5 (98)	4.4(160)	10.5 (76)	442	17.0 (86)	12.9 (66)	666	3.8 (11)
1959	7	* *	* *	80	* *	* *	* *	44	* *	* *	56	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 31
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
2150	254	9.1 (70)	8.7 (86)	1,122	7.6(165)	1.0 (12)	12.7(145)	546	15.9 (58)	17.2(142)	702 3.0 (5)
2151	95	* *	* *	408	5.6(103)	2.0 (63)	10.3 (71)	195	19.5(132)	12.3 (51)	265 3.4 (8)
2251	165	7.3 (30)	7.3 (63)	669	5.8(112)	1.0 (12)	11.2 (95)	311	16.4 (69)	12.9 (66)	458 6.1 (33)
2252	75	* *	* *	356	7.6(165)	2.2 (78)	12.6(140)	196	27.0(158)	14.3 (97)	248 9.7 (66)
2551	12	* *	* *	250	8.0(176)	2.8(114)	11.2 (95)	104	18.3(117)	12.5 (58)	170 18.8(133)
2552	6	* *	* *	9	* *	* *	* *	1	* *	* *	2 * *
2554	45	* *	* *	230	7.4(161)	1.3 (16)	18.3(180)	124	22.6(152)	19.4(151)	162 10.5 (74)
2555	2,808	11.2 (97)	4.9 (16)	10,611	6.8(154)	2.6(100)	10.1 (62)	4,339	15.7 (55)	12.4 (55)	6,538 17.3(128)
3001	328	12.8 (109)	4.9 (16)	1,591	5.3 (85)	2.6(100)	14.2(167)	828	18.0(109)	15.2(117)	1,120 9.1 (62)
3002	83	* *	* *	446	5.2 (79)	1.3 (16)	14.6(172)	241	21.6(147)	14.1 (89)	328 16.8(125)
3003	2	* *	* *	4	* *	* *	* *	1	* *	* *	1 * *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 32
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
3004	41	* *	* *	271	5.2 (79)	1.8 (48)	21.0(183)	159	24.5(155)	21.4(157)	214 10.3 (70)
3007	3	* *	* *	54	* *	* *	* *	37	* *	* *	49 * *
3009	3	* *	* *	13	* *	* *	* *	9	* *	* *	11 * *
3201	196	9.7 (80)	10.7 (105)	911	5.9(120)	3.0(125)	14.6(172)	412	15.8 (57)	16.5(135)	609 8.5 (57)
3301	1,389	8.2 (48)	7.2 (57)	5,392	5.4 (90)	3.5(144)	12.0(124)	2,336	17.4 (98)	13.5 (79)	3,474 12.8 (96)
3302	61	* *	* *	450	6.7(150)	2.2 (78)	14.0(163)	238	18.1(113)	17.2(142)	331 14.5 (115)
3304	58	* *	* *	272	3.3 (12)	4.4(160)	7.7 (14)	140	14.3 (34)	11.4 (30)	188 8.5 (57)
3305	65	* *	* *	281	3.9 (26)	4.3(157)	7.1 (10)	135	16.3 (66)	7.4 (6)	189 9.0 (61)
3306	100	5.0 (3)	14.0 (115)	461	4.8 (57)	2.6(100)	8.7 (27)	229	18.8(123)	10.9 (22)	304 10.9 (78)
3308	16	* *	* *	115	4.3 (39)	0.9 (10)	13.9(161)	55	* *	* *	81 * *
3309	19	* *	* *	95	* *	* *	* *	52	* *	* *	64 * *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 33
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
3550	30	* *	* *	147	3.4 (14)	0.0 (1)	23.1(184)	80	* *	* *	80 * *
3551	9	* *	* *	44	* *	* *	* *	25	* *	* *	25 * *
3552	15	* *	* *	87	* *	* *	* *	46	* *	* *	50 * *
3701	422	8.3 (52)	6.9 (50)	1,821	4.8 (57)	3.4(141)	9.3 (42)	841	20.6(139)	12.1 (43)	1,247 7.8 (51)
3805	1,131	10.1 (87)	7.9 (75)	4,915	4.6 (52)	3.3(137)	9.4 (43)	2,277	17.2 (92)	12.9 (66)	2,934 13.7 (107)
3950	605	6.3 (13)	10.1 (101)	2,941	4.5 (47)	3.8(151)	11.2 (95)	1,425	18.7(122)	12.4 (55)	1,971 25.1 (151)
3952	1	* *	* *	2	* *	* *	* *	1	* *	* *	1 * *
3954	19	* *	* *	103	4.9 (67)	2.9(119)	11.7(116)	61	* *	* *	64 * *
4001	442	14.3 (113)	6.3 (37)	1,923	5.7(106)	3.8(151)	12.0(124)	915	16.4 (69)	12.7 (62)	1,305 36.9 (163)
4002	398	9.5 (77)	6.0 (31)	1,821	4.8 (57)	3.2(132)	12.6(140)	873	14.3 (34)	13.9 (88)	1,279 22.0 (144)
4103	36	* *	* *	226	5.3 (85)	4.0(154)	11.9(118)	131	13.7 (30)	12.2 (47)	151 27.2 (153)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 34
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature		Under- weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4106	787	11.3 (99)	6.5 (43)	3,388	6.6(146)	1.4 (21)	11.9(118)	1,627	16.7 (78)	14.5(103)	2,283	14.1 (110)
4107	1,251	9.8 (81)	8.2 (79)	5,210	6.4(141)	1.8 (48)	10.1 (62)	2,424	17.2 (92)	11.9 (36)	3,474	15.8 (118)
4109	390	8.7 (61)	8.5 (83)	1,940	4.8 (57)	2.4 (87)	8.6 (25)	996	17.5(102)	11.0 (23)	1,328	6.2 (34)
4110	85	* *	* *	908	5.4 (90)	1.3 (16)	14.9(174)	511	20.2(135)	15.9(126)	667	10.0 (68)
4112	195	3.1 (2)	7.7 (72)	1,060	5.8(112)	3.8(151)	17.4(179)	582	19.2(129)	20.6(155)	863	11.8 (89)
4113	265	3.0 (1)	12.5(113)	1,170	3.0 (4)	1.8 (48)	13.8(158)	590	21.4(146)	20.0(153)	785	7.3 (47)
4120	751	8.7 (61)	7.5 (69)	3,412	6.6(146)	1.7 (38)	10.7 (85)	1,612	16.1 (61)	11.8 (35)	2,271	11.1 (81)
4160	625	11.0 (92)	8.3 (81)	2,910	6.8(154)	2.0 (63)	10.8 (87)	1,314	15.9 (58)	12.2 (47)	1,896	12.2 (90)
4351	183	6.6 (17)	9.8 (99)	968	5.8(112)	1.7 (38)	9.6 (49)	481	21.6(147)	11.2 (27)	683	14.1 (110)
4352	64	* *	* *	296	5.4 (90)	2.7(109)	12.2(131)	167	18.0(109)	15.0(112)	215	12.6 (94)
4353	126	11.9(102)	6.3 (37)	779	5.3 (85)	3.2(132)	11.9(118)	395	21.8(149)	12.7 (62)	584	11.3 (84)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 35
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4354	275	5.5 (7)	9.1 (91)	1,221	5.5 (98)	1.6 (34)	9.2 (40)	549	17.1 (88)	12.8 (64)	832	5.9 (31)
4355	90	* *	* *	498	3.4 (14)	2.4 (87)	12.4(136)	255	19.2(129)	17.3(144)	357	10.9 (78)
4356	182	7.1 (27)	3.8 (8)	900	5.7(106)	0.8 (7)	12.0(124)	454	21.8(149)	15.0(112)	634	12.8 (96)
4358	43	* *	* *	327	5.8(112)	0.3 (5)	20.8(182)	191	24.6(156)	30.9(159)	255	15.7 (117)
4359	201	6.5 (15)	14.4 (117)	958	5.4 (90)	1.7 (38)	13.3(153)	514	20.6(139)	13.8 (87)	677	10.5 (74)
4360	67	* *	* *	304	4.6 (52)	2.0 (63)	8.9 (32)	155	16.1 (61)	12.3 (51)	208	9.1 (62)
4361	68	* *	* *	357	7.6(165)	1.4 (21)	12.6(140)	183	24.6(156)	20.2(154)	245	13.1 (100)
4362	101	7.9 (36)	9.9(100)	518	5.6(103)	0.8 (7)	9.5 (47)	269	14.9 (41)	13.4 (77)	361	12.5 (92)
4363	320	6.6 (17)	9.4 (95)	1,422	5.1 (74)	1.7 (38)	8.9 (32)	666	16.2 (65)	12.6 (60)	992	10.0 (68)
4401	500	8.2 (48)	7.8 (73)	1,973	5.9(120)	2.8(114)	10.9 (90)	888	16.0 (60)	14.8(111)	1,294	7.9 (52)
4501	330	7.9 (36)	7.9 (75)	1,644	5.4 (90)	1.5 (27)	9.2 (40)	803	13.9 (31)	12.1 (43)	1,174	6.5 (38)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 36
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)													
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature			Under- weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)		Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4502	1	* *	* *	3	* *	* *	* *		1	* *	* *	2	* *
4591	0	* *	* *	1	* *	* *	* *		1	* *	* *	1	* *
4601	615	7.5 (33)	6.3 (37)	2,272	5.5 (98)	2.9(119)	12.3(133)		991	21.2(144)	15.6(123)	1,220	6.9 (41)
4701	516	6.4 (14)	10.9(108)	2,021	5.8(112)	1.6 (34)	10.2 (67)		883	15.7 (55)	14.6(105)	1,351	5.6 (29)
5050	605	12.4 (105)	7.3 (63)	2,429	5.6(103)	1.8 (48)	11.6(112)		1,024	13.0 (24)	12.1 (43)	1,579	13.9(109)
5051	1,027	9.3 (74)	6.5 (43)	3,881	6.2(129)	2.2 (78)	11.3(101)		1,670	12.8 (22)	11.6 (33)	2,492	14.2 (112)
5052	970	9.9 (84)	5.4 (24)	3,841	5.4 (90)	1.5 (27)	11.6(112)		1,630	17.4 (98)	13.7 (84)	2,534	12.9 (98)
5101	835	8.6 (59)	5.5 (27)	3,272	6.0(123)	2.1 (73)	10.9 (90)		1,414	17.5(102)	14.1 (89)	2,093	8.8 (59)
5202	237	6.8 (22)	8.0 (77)	942	6.3(139)	2.4 (87)	13.1(148)		416	20.2(135)	16.3(130)	569	7.0 (43)
5203	40	* *	* *	235	6.0(123)	1.7 (38)	11.9(118)		125	15.2 (48)	14.4(100)	168	14.9(116)
5204	41	* *	* *	178	6.7(150)	1.1 (14)	10.1 (62)		90	* *	* *	111	12.6 (94)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 37
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
5601	439	9.6 (79)	9.6 (98)	2,013	5.3 (85)	1.7 (38)	12.9(146)	935	15.1 (46)	14.4(100)	1,257	8.2 (56)
5801	697	10.5 (91)	6.9 (50)	2,782	7.6(165)	1.7 (38)	11.7(116)	1,223	17.8(106)	15.1(116)	1,861	8.1 (55)
5803	76	* *	* *	427	3.7 (21)	2.1 (73)	11.2 (95)	197	12.7 (19)	13.7 (84)	304	10.5 (74)
5804	19	* *	* *	167	4.2 (34)	4.2(156)	13.2(150)	91	* *	* *	129	7.0 (43)
6101	22	* *	* *	125	3.2 (10)	5.6(170)	6.4 (5)	70	* *	* *	92	* *
6102	69	* *	* *	393	6.1(127)	4.6(165)	11.2 (95)	214	12.6 (18)	13.6 (82)	278	13.3(104)
6103	1,300	9.8 (81)	6.1 (33)	5,689	5.2 (79)	3.5(144)	10.0 (58)	2,744	17.2 (92)	12.0 (41)	3,865	18.7 (132)
6104	262	9.2 (72)	5.0 (19)	677	4.3 (39)	5.6(170)	10.0 (58)	245	20.0(134)	18.4(150)	362	14.4 (113)
6105	20	* *	* *	130	6.2(129)	0.8 (7)	8.5 (23)	66	* *	* *	98	* *
6106	1	* *	* *	4	* *	* *	* *	2	* *	* *	3	* *
6301	1,800	9.8 (81)	5.3 (21)	6,755	6.2(129)	2.0 (63)	8.6 (25)	2,764	16.1 (61)	12.9 (66)	4,309	13.2(102)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 38
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
6312	1,095	11.0 (92)	7.0 (53)	4,159	7.1(160)	2.8(114)	8.2 (18)	1,693	14.7 (38)	10.5 (18)	2,681	17.9 (130)
6316	627	10.2 (88)	7.2 (57)	2,311	5.1 (74)	2.1 (73)	9.0 (38)	873	14.0 (32)	14.1 (89)	1,480	11.2 (83)
6320	60	* *	* *	383	3.1 (8)	2.9(119)	10.4 (73)	210	12.4 (13)	11.9 (36)	262	4.6 (18)
6325	44	* *	* *	253	7.9(175)	3.6(147)	9.9 (55)	128	23.4(154)	14.1 (89)	169	4.1 (16)
6330	143	16.1 (116)	6.3 (37)	626	7.5(163)	2.6(100)	9.4 (43)	315	12.4 (13)	8.6 (9)	399	18.8(133)
6335	175	8.6 (59)	10.3(104)	966	3.5 (17)	3.5(144)	8.4 (22)	487	15.4 (50)	10.3 (17)	649	7.9 (52)
6340	47	* *	* *	207	4.3 (39)	1.4 (21)	5.3 (2)	115	11.3 (8)	8.7 (10)	135	3.7 (10)
6345	4	* *	* *	45	* *	* *	* *	23	* *	* *	28	* *
6350	72	* *	* *	421	4.0 (29)	3.3(137)	11.6(112)	191	11.0 (7)	15.2(117)	278	11.5 (85)
7304	67	* *	* *	356	4.2 (34)	2.0 (63)	13.8(158)	187	23.0(153)	17.6(146)	253	3.2 (7)
7306	955	11.4 (100)	7.4 (66)	3,927	5.1 (74)	2.6(100)	12.0(124)	1,816	16.1 (61)	16.3(130)	2,687	13.1 (100)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 39
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
7308	58	* *	* *	277	4.0 (29)	2.5 (94)	13.4(154)	132	13.6 (29)	16.7(138)	189 5.3 (25)
7310	23	* *	* *	71	* *	* *	* *	26	* *	* *	45 * *
7401	921	6.9 (23)	6.2 (34)	3,926	5.7(106)	1.9 (59)	7.9 (15)	1,777	16.5 (73)	11.9 (36)	2,604 6.6 (40)
7601	244	7.0 (26)	6.6 (45)	1,090	3.9 (26)	2.4 (87)	9.4 (43)	497	16.7 (78)	13.3 (75)	755 6.5 (38)
7850	388	8.5 (55)	4.9 (16)	1,700	4.9 (67)	4.9(167)	11.9(118)	828	18.2(115)	16.3(130)	1,165 4.3 (17)
7852	41	* *	* *	160	3.1 (8)	5.6(170)	8.8 (30)	73	* *	* *	112 6.3 (36)
8101	434	22.1 (117)	4.4 (12)	713	7.4(161)	11.1(184)	8.7 (27)	205	12.2 (12)	11.2 (27)	307 32.6(161)
8102	31	* *	* *	134	8.2(179)	6.0(175)	14.2(167)	75	* *	* *	99 * *
8105	103	8.7 (61)	8.7 (86)	216	10.6(184)	1.4 (21)	12.5(137)	67	* *	* *	94 * *
8106	62	* *	* *	164	4.9 (67)	3.7(150)	14.0(163)	86	* *	* *	109 30.3(155)
8150	14	* *	* *	90	* *	* *	* *	53	* *	* *	66 * *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 40
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
8151	1,049	9.2 (72)	7.1 (55)	4,307	6.2(129)	1.7 (38)	11.1 (94)	1,843	17.4 (98)	15.0(112)	2,821 19.8 (137)
8152	143	12.6 (107)	4.2 (11)	393	6.6(146)	4.1(155)	10.2 (67)	112	12.5 (15)	17.9(149)	204 45.1 (166)
8153	13	* *	* *	48	* *	* *	* *	28	* *	* *	33 * *
8154	121	5.0 (3)	9.1 (91)	530	3.0 (4)	3.6(147)	6.8 (7)	257	15.6 (52)	9.3 (11)	350 6.9 (41)
8155	26	* *	* *	135	2.2 (2)	3.0(125)	9.6 (49)	73	* *	* *	98 * *
8156	5	* *	* *	31	* *	* *	* *	19	* *	* *	22 * *
8157	2	* *	* *	33	* *	* *	* *	25	* *	* *	26 * *
8201	959	8.1 (40)	6.3 (37)	3,275	5.8(112)	2.4 (87)	9.4 (43)	1,071	19.1(127)	12.3 (51)	1,758 3.0 (5)
8202	1,159	8.1 (40)	7.2 (57)	5,250	4.9 (67)	1.7 (38)	10.6 (79)	2,568	17.8(106)	12.4 (55)	3,439 25.3 (152)
8203	135	8.1 (40)	5.9 (30)	548	6.8(154)	0.5 (6)	10.6 (79)	218	19.3(131)	11.9 (36)	324 13.3 (104)
8250	390	11.0 (92)	6.4 (42)	1,334	5.0 (72)	2.9(119)	8.3 (21)	465	11.8 (9)	8.2 (8)	677 17.7 (129)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 41
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
8252	45	* *	* *	125	10.4(182)	8.0(181)	7.2 (11)	63	* *	* *	75 * *
8255	1,177	8.5 (55)	8.2 (79)	3,833	7.8(173)	1.1 (14)	10.1 (62)	1,323	20.2(135)	11.0 (23)	2,106 13.2 (102)
8257	114	8.8 (65)	5.3 (21)	771	5.8(112)	4.8(166)	10.6 (79)	376	13.0 (24)	14.6(105)	500 9.4 (65)
8258	387	8.5 (55)	5.7 (29)	1,771	6.9(157)	1.5 (27)	13.4(154)	774	20.7(141)	11.9 (36)	1,046 16.1 (121)
8260	307	9.4 (76)	2.6 (2)	1,208	5.7(106)	3.2(132)	8.2 (18)	549	13.1 (26)	11.3 (29)	812 16.6 (123)
8901	431	13.7 (112)	2.6 (2)	1,756	5.8(112)	5.2(169)	9.6 (49)	764	14.4 (37)	12.0 (41)	1,127 31.1 (159)
8902	359	11.1 (96)	2.8 (4)	1,226	5.1 (74)	7.6(179)	9.1 (39)	513	12.9 (23)	9.7 (13)	760 30.5 (156)
8903	474	14.8 (115)	3.6 (6)	2,036	6.9(157)	4.5(163)	12.6(140)	922	16.7 (78)	13.1 (72)	1,343 39.2 (165)
8904	487	12.3 (103)	6.2 (34)	1,814	4.5 (47)	5.8(174)	8.5 (23)	742	15.0 (44)	9.7 (13)	1,161 32.2 (160)
8905	1,004	8.5 (55)	5.0 (19)	4,782	4.6 (52)	6.7(176)	13.2(150)	2,373	16.9 (84)	13.5 (79)	3,422 20.9 (141)
9004	1	* *	* *	1	* *	* *	* *	0	* *	* *	0 * *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 42
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
9010	6	* *	* *	3	* *	* *	* *	0	* *	* *	0 * *
9012	15	* *	* *	89	* *	* *	* *	50	* *	* *	20 * *
9013	47	* *	* *	166	4.8 (57)	3.6(147)	5.4 (3)	64	* *	* *	20 * *
9015	19	* *	* *	54	* *	* *	* *	31	* *	* *	22 * *
9016	69	* *	* *	290	6.2(129)	3.1(130)	10.7 (85)	121	17.4 (98)	11.6 (33)	149 45.6 (167)
9017	26	* *	* *	114	10.5(183)	0.0 (1)	18.4(181)	58	* *	* *	22 * *
9018	51	* *	* *	194	8.2(179)	3.1(130)	10.8 (87)	81	* *	* *	9 * *
9019	16	* *	* *	70	* *	* *	* *	34	* *	* *	9 * *
9021	34	* *	* *	160	3.8 (23)	4.4(160)	13.8(158)	77	* *	* *	16 * *
9040	0	* *	* *	1	* *	* *	* *	1	* *	* *	0 * *
9041	9	* *	* *	100	3.0 (4)	2.0 (63)	6.0 (4)	76	* *	* *	23 * *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 43
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators												(3,4)	
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature		Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)	
9045	555	11.2 (97)	7.4 (66)	2,539	5.9(120)	1.3 (16)	13.4(154)	1,199	18.0(109)	15.7(124)	1,701	24.2 (148)	
9050	471	10.0 (86)	5.3 (21)	2,391	6.2(129)	2.6(100)	12.3(133)	1,123	15.5 (51)	13.7 (84)	1,740	17.1 (127)	
9051	532	7.9 (36)	8.5 (83)	2,604	6.5(143)	2.5 (94)	16.4(178)	1,350	17.3 (97)	17.8(148)	1,866	16.7 (124)	
9053	936	14.7 (114)	4.1 (9)	3,032	6.4(141)	2.8(114)	8.1 (17)	1,184	16.7 (78)	11.1 (26)	1,880	24.8 (150)	
9055	11	* *	* *	9	* *	* *	* *	1	* *	* *	1	* *	
9058	252	9.9 (84)	6.0 (31)	1,332	3.5 (17)	3.0(125)	9.7 (53)	680	11.8 (9)	10.6 (19)	949	22.2 (146)	
9066	626	12.9 (111)	5.4 (24)	2,669	4.8 (57)	5.6(170)	7.0 (9)	1,142	8.5 (4)	6.8 (5)	1,819	9.8 (67)	
9068	422	9.5 (77)	3.6 (6)	1,879	6.0(123)	9.5(182)	7.5 (12)	812	8.0 (3)	5.3 (2)	1,284	24.2 (148)	
9069	695	12.8 (109)	4.7 (13)	2,564	7.7(172)	2.3 (83)	8.8 (30)	922	14.3 (34)	11.0 (23)	1,574	20.9 (141)	
9070	514	10.3 (89)	2.5 (1)	2,355	8.0(176)	7.9(180)	11.4(107)	1,011	7.3 (1)	6.1 (3)	1,547	27.9 (154)	
9072	516	11.0 (92)	3.3 (5)	1,987	7.8(173)	2.2 (78)	10.0 (58)	809	12.5 (15)	9.6 (12)	1,274	16.0 (120)	

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 44
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9074	370	12.7 (108)	5.4 (24)	1,528	6.2(129)	3.4(141)	10.1 (62)	659	12.7 (19)	10.6 (19)	1,020	12.9 (98)
9080	973	12.3 (103)	4.1 (9)	4,086	5.5 (98)	2.3 (83)	8.9 (32)	1,857	15.6 (52)	12.2 (47)	2,645	33.9 (162)
Unknown Clinics	124	5.6 (8)	4.8 (14)	467	5.4 (90)	3.4(141)	15.6(175)	240	20.4 (138)	22.5(158)	329	12.2 (90)
Michigan	59,582	9.4	6.9	254,034	5.7	2.8	10.9	116,006	16.6	13.3	166,748	15.0
Nation (Prior Year)	1,851,546	9.0	6.3	8,057,124	6.3	3.6	12.4	3,572,939	16.1	14.4	5,391,027	14.6

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 45
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)									
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number	% (Rank)
0101	39	* *	28 * *	29 * *	18 * *	26 * *	104 85.6 (91)	197	7.1 (114)
0102	175	70.3 (27)	115 17.4 (63)	110 8.2 (62)	85 * *	108 3.7 (72)	299 87.3 (73)	652	8.9 (145)
0103	167	59.9 (71)	111 14.4 (79)	97 * *	95 * *	108 4.6 (61)	327 84.1(101)	656	11.0 (165)
0104	47	* *	44 * *	35 * *	31 * *	43 * *	98 * *	210	14.3 (177)
0250	48	* *	36 * *	48 * *	24 * *	33 * *	125 91.2 (38)	252	7.5 (123)
0251	30	* *	30 * *	28 * *	21 * *	29 * *	101 82.2(117)	186	7.0 (108)
0252	33	* *	27 * *	48 * *	15 * *	27 * *	127 85.0 (95)	220	9.5 (150)
0253	58	* *	55 * *	37 * *	51 * *	54 * *	161 87.0 (78)	294	12.6 (170)
0401	177	51.4 (94)	157 19.1 (50)	157 10.8 (26)	126 23.0 (11)	155 8.4 (31)	409 91.9 (31)	817	5.1 (68)
0402	130	61.5 (66)	89 * *	105 9.5 (44)	63 * *	81 * *	321 86.9 (79)	613	8.5 (136)
0403	41	* *	34 * *	40 * *	24 * *	32 * *	129 83.7(103)	228	7.9 (128)
0404	43	* *	34 * *	46 * *	36 * *	34 * *	124 84.7 (97)	240	21.7 (184)
0501	210	76.7 (10)	167 29.3 (8)	155 11.0 (24)	126 27.8 (6)	151 14.6 (5)	456 90.8 (41)	912	12.9 (173)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 46
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)								
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)	(Rank)
0502	0	* *	2	* *	2	* *	2	* *
0503	60	* *	52	* *	49	* *	34	* *
0504	197	80.2 (6)	158	33.5 (5)	137	19.7 (3)	135	34.8 (1)
0505	144	81.9 (3)	97	* *	89	* *	147	10.9 (15)
0506	67	* *	50	* *	52	* *	135	34.8 (1)
0650	363	76.3 (11)	295	23.7 (28)	277	9.7 (40)	93	* *
0651	231	62.3 (64)	172	12.8 (87)	193	6.7 (78)	94	* *
0652	104	60.6 (67)	95	* *	89	* *	315	90.8 (41)
0653	138	65.9 (51)	125	27.2 (14)	148	8.8 (52)	615	5.7 (78)
0654	136	68.4 (37)	112	17.9 (58)	117	9.4 (45)	196	85.2 (92)
0655	148	73.6 (16)	123	24.4 (25)	93	* *	338	11.2 (167)
0657	24	* *	16	* *	17	* *	796	90.1 (55)
0701	137	86.9 (1)	113	38.9 (2)	106	19.8 (2)	1,590	6.8 (103)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 47
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)									
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number	% (Rank)
0702	29	* *	22 * *	29 * *	13 * *	21 * *	72 * *	144	2.8 (20)
0703	74	* *	64 * *	51 * *	45 * *	60 * *	183 90.7 (45)	356	7.0 (108)
0704	28	* *	12 * *	17 * *	12 * *	12 * *	86 * *	143	10.5 (161)
0705	53	* *	53 * *	34 * *	38 * *	52 * *	150 94.7 (11)	284	1.8 (13)
0850	408	74.5 (13)	376 19.9 (46)	335 7.8 (67)	276 19.2 (22)	353 9.1 (26)	847 88.1 (66)	1,766	4.2 (51)
0851	284	68.3 (38)	284 19.4 (49)	267 10.5 (29)	189 21.7 (14)	277 9.0 (28)	768 90.6 (48)	1,459	3.6 (31)
0901	609	58.1 (77)	495 13.5 (83)	509 5.1 (84)	445 12.6 (49)	482 3.3 (77)	1,377 94.2 (13)	2,766	7.0 (108)
0902	24	* *	22 * *	31 * *	19 * *	20 * *	68 * *	143	5.6 (75)
1051	556	81.8 (4)	353 39.9 (1)	368 14.9 (6)	219 32.9 (2)	331 15.4 (2)	1,023 91.9 (31)	2,151	2.8 (20)
1101	713	46.1(101)	429 12.4 (89)	495 8.3 (59)	311 9.6 (62)	422 4.3 (66)	1,419 73.8(144)	2,895	7.2 (117)
1102	334	56.0 (85)	229 17.9 (58)	268 9.7 (40)	171 16.4 (32)	224 7.6 (40)	637 81.0(122)	1,381	9.4 (149)
1106	41	* *	28 * *	32 * *	24 * *	26 * *	78 * *	174	2.9 (22)
1250	347	58.8 (74)	291 21.0 (42)	236 9.7 (40)	250 16.8 (31)	282 9.9 (23)	929 82.3(116)	1,753	7.2 (117)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 48
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)		Number % (Rank)	
1251	312	69.9 (32)	240 23.8 (26)	199 10.1 (32)	219 18.3 (27)	221 10.0 (21)	721 91.4 (36)		1,447 6.9 (106)	
1253	547	65.8 (53)	436 25.0 (23)	372 9.4 (45)	350 14.9 (36)	412 13.3 (8)	1,246 86.8 (82)		2,440 4.1 (47)	
1350	840	57.5 (78)	631 12.5 (88)	612 3.6 (93)	527 9.9 (60)	584 3.9 (69)	1,915 89.5 (59)		3,842 4.6 (61)	
1351	45	* *	31 * *	30 * *	30 * *	31 * *	125 93.6 (17)		218 3.7 (38)	
1352	162	53.7 (89)	122 10.7 (95)	108 3.7 (92)	91 * *	119 2.5 (87)	394 87.6 (70)		734 10.2 (155)	
1401	66	* *	48 * *	54 * *	36 * *	45 * *	189 92.1 (27)		357 3.6 (31)	
1402	48	* *	34 * *	47 * *	26 * *	33 * *	150 88.0 (67)		272 5.5 (72)	
1403	11	* *	10 * *	11 * *	10 * *	10 * *	29 * *		58 * *	
1404	4	* *	5 * *	2 * *	4 * *	5 * *	18 * *		29 * *	
1501	352	70.2 (30)	282 23.8 (26)	328 12.2 (19)	248 21.4 (16)	258 7.8 (39)	1,028 83.9(102)		1,880 6.4 (91)	
1503	402	66.4 (48)	339 20.1 (44)	345 10.1 (32)	268 13.1 (44)	313 8.3 (33)	1,100 86.9 (79)		2,049 6.3 (89)	
1504	875	72.7 (20)	591 26.1 (19)	724 11.5 (22)	434 23.7 (9)	524 8.6 (30)	1,996 85.8 (89)		3,923 3.4 (25)	
1505	309	77.3 (9)	226 34.1 (4)	245 13.1 (15)	178 28.1 (5)	208 15.4 (2)	794 92.4 (26)		1,471 1.4 (12)	

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 49
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
1506	226	72.6 (21)	171	24.6 (24)	190	12.6 (17)	135	18.5 (26)	162	9.3 (25)	573	92.8 (22)	1,084	4.2 (51)
1507	230	73.5 (18)	157	25.5 (21)	172	9.9 (35)	120	27.5 (7)	155	14.8 (4)	529	91.9 (31)	1,035	2.3 (18)
1508	37	* *	29	* *	47	* *	13	* *	22	* *	124	92.7 (24)	229	0.9 (9)
1509	185	70.3 (27)	163	27.0 (16)	173	13.9 (10)	111	11.7 (53)	149	6.7 (45)	600	86.0 (85)	1,046	5.0 (66)
1511	301	74.4 (14)	225	21.3 (40)	183	10.9 (25)	200	21.0 (17)	217	5.5 (54)	559	92.8 (22)	1,151	4.1 (47)
1512	127	63.0 (59)	118	25.4 (22)	93	* *	95	* *	116	11.2 (11)	359	96.4 (4)	665	3.6 (31)
1701	259	68.7 (36)	177	31.1 (7)	148	14.9 (6)	166	32.5 (3)	165	21.8 (1)	581	86.9 (79)	1,170	6.8 (103)
1950	240	71.7 (24)	205	15.6 (71)	161	8.1 (64)	169	14.8 (38)	201	8.0 (35)	471	90.7 (45)	1,017	7.5 (123)
1952	45	* *	43	* *	42	* *	26	* *	39	* *	127	90.6 (48)	247	7.7 (127)
1953	88	* *	88	* *	75	* *	65	* *	84	* *	272	87.1 (76)	509	6.7 (97)
1954	212	70.3 (27)	162	17.3 (65)	130	8.5 (54)	143	18.9 (23)	161	8.7 (29)	484	87.2 (75)	956	8.4 (133)
1955	53	* *	39	* *	35	* *	31	* *	35	* *	144	83.3(107)	266	10.2 (155)
1957	219	81.7 (5)	198	27.8 (11)	151	10.6 (28)	167	28.7 (4)	189	10.1 (19)	475	87.4 (72)	986	3.7 (38)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 50
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)						Exclusive Breastfeeding (7)		≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
1959	7	* *	22	* *	14	* *	10	* *	21	* *	43	* *
2150	254	65.0 (55)	184	20.1 (44)	168	8.3 (59)	159	17.6 (30)	177	6.2 (50)	595	93.8 (16)
2151	95	* *	70	* *	70	* *	58	* *	66	* *	210	90.5 (51)
2251	165	66.1 (49)	121	22.3 (35)	113	8.0 (66)	92	* *	117	9.4 (24)	336	89.0 (62)
2252	75	* *	47	* *	41	* *	39	* *	43	* *	207	91.3 (37)
2551	12	* *	65	* *	47	* *	33	* *	64	* *	115	80.0(126)
2552	6	* *	1	* *	1	* *	1	* *	1	* *	1	* *
2554	45	* *	39	* *	38	* *	33	* *	39	* *	128	94.5 (12)
2555	2,809	52.3 (91)	2,375	11.5 (94)	1,556	5.8 (80)	1,959	8.1 (72)	2,257	3.2 (78)	4,582	78.8(132)
3001	327	53.8 (87)	267	6.0 (106)	219	1.4 (104)	224	4.5 (81)	261	1.1(106)	894	83.0(110)
3002	84	* *	65	* *	63	* *	52	* *	64	* *	255	95.3 (8)
3003	2	* *	0	* *	0	* *	0	* *	0	* *	1	* *
3004	42	* *	36	* *	38	* *	31	* *	34	* *	179	72.6(147)
											283	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years ⁽³⁾**

(2)

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
- (8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 52
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
3701	422	70.1 (31)	362	16.9 (68)	357	9.8 (38)	286	13.3 (42)	338	8.3 (33)
3805	1,136	63.0 (59)	1,070	14.5 (77)	900	8.1 (64)	758	12.5 (50)	1,005	6.2 (50)
3950	611	66.9 (45)	498	29.3 (8)	517	14.7 (8)	396	21.7 (14)	470	11.1 (13)
3952	1	* *	1	* *	0	* *	1	* *	1	* *
3954	19	* *	10	* *	10	* *	10	* *	66	* *
4001	462	60.4 (68)	303	18.2 (55)	285	7.0 (75)	271	14.8 (38)	278	6.8 (44)
4002	402	65.9 (51)	322	17.4 (63)	285	9.1 (50)	297	13.1 (44)	310	3.2 (78)
4103	37	* *	26	* *	30	* *	31	* *	25	* *
4106	793	62.9 (62)	583	20.8 (43)	534	8.4 (56)	477	8.8 (67)	530	5.1 (58)
4107	1,257	67.5 (42)	1,038	22.9 (31)	793	9.8 (38)	828	18.8 (24)	970	10.8 (16)
4109	393	66.9 (45)	343	22.7 (34)	303	14.2 (9)	275	19.3 (21)	323	10.2 (18)
4110	85	* *	148	18.2 (55)	130	7.7 (69)	95	* *	132	4.5 (62)
4112	195	28.2(116)	153	16.3 (70)	161	13.7 (12)	133	4.5 (81)	149	2.0 (91)
	625	96.2 (6)								

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 53
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
4113	265	67.5 (42)	163	22.1 (37)	135	12.6 (17)	167	9.0 (66)	161	3.1 (81)
4120	754	66.0 (50)	641	18.1 (57)	525	8.4 (56)	498	9.8 (61)	599	5.7 (53)
4160	633	71.6 (25)	587	27.6 (12)	483	17.0 (4)	437	19.5 (19)	531	13.7 (7)
4351	185	61.6 (65)	164	22.0 (39)	161	9.9 (35)	133	15.8 (34)	161	11.8 (10)
4352	64	* *	40	* *	37	* *	31	* *	38	* *
4353	127	70.9 (26)	153	22.2 (36)	161	9.3 (48)	114	20.2 (18)	149	10.1 (19)
4354	276	72.1 (22)	242	18.6 (53)	208	5.8 (80)	221	22.2 (12)	241	12.0 (9)
4355	90	* *	82	* *	88	* *	65	* *	78	* *
4356	182	67.0 (44)	155	27.1 (15)	158	13.9 (10)	125	23.2 (10)	150	8.0 (35)
4358	43	* *	39	* *	60	* *	31	* *	38	* *
4359	202	79.2 (7)	117	23.1 (29)	147	13.6 (14)	89	* *	108	2.8 (83)
4360	67	* *	52	* *	49	* *	51	* *	51	* *
4361	68	* *	51	* *	61	* *	29	* *	50	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 54
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)						Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
4362	101	79.2 (7)	72	* *	90	* *	51	* *	72	* *	288	89.2 (61)
4363	320	73.1 (19)	272	19.1 (50)	238	9.7 (40)	219	13.2 (43)	268	4.5 (62)	708	87.9 (68)
4401	502	66.7 (47)	383	17.2 (67)	349	8.3 (59)	315	14.9 (36)	368	7.3 (42)	953	93.4 (19)
4501	330	51.8 (93)	290	13.1 (84)	246	8.5 (54)	231	10.0 (59)	284	1.8 (96)	864	91.8 (34)
4502	1	* *	1	* *	1	* *	1	* *	1	* *	1	* *
4591	0	* *	0	* *	0	* *	0	* *	0	* *	1	* *
4601	623	69.7 (34)	415	19.5 (48)	355	10.4 (30)	343	15.2 (35)	392	7.1 (43)	1,067	76.9(139)
4701	514	69.8 (33)	355	27.3 (13)	391	10.0 (34)	262	21.8 (13)	340	10.0 (21)	946	92.1 (27)
5050	608	59.7 (72)	429	14.0 (82)	469	5.3 (83)	387	8.3 (70)	407	3.9 (69)	1,077	82.5(114)
5051	1,026	57.5 (78)	666	14.6 (75)	671	7.2 (71)	637	8.8 (67)	633	6.3 (49)	1,788	78.7(133)
5052	969	53.8 (87)	716	12.2 (90)	702	5.7 (82)	618	6.5 (75)	675	4.0 (67)	1,724	78.4(134)
5101	844	68.1 (40)	602	22.8 (33)	605	9.9 (35)	538	14.3 (41)	554	6.7 (45)	1,507	86.5 (83)
5202	237	73.8 (15)	186	23.1 (29)	176	10.8 (26)	154	19.5 (19)	177	7.9 (37)	459	90.0 (56)
											969	4.5 (59)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 55
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)						Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
5203	40	* *	36	* *	51	* *	25	* *	36	* *	133	82.7 (111)
5204	41	* *	30	* *	28	* *	22	* *	28	* *	101	87.1 (76)
5601	439	73.6 (16)	383	26.4 (17)	323	8.7 (53)	325	24.9 (8)	362	9.1 (26)	998	90.7 (45)
5801	700	63.3 (58)	554	14.6 (75)	497	7.0 (75)	452	11.7 (53)	531	5.3 (56)	1,322	82.5 (114)
5803	76	* *	89	* *	80	* *	70	* *	85	* *	219	91.8 (34)
5804	19	* *	32	* *	26	* *	20	* *	31	* *	97	* *
6101	22	* *	19	* *	19	* *	17	* *	17	* *	73	* *
6102	69	* *	48	* *	55	* *	54	* *	47	* *	238	86.1 (84)
6103	1,302	56.2 (83)	878	15.4 (72)	915	7.2 (71)	825	13.0 (47)	844	4.7 (60)	2,932	85.1 (93)
6104	263	42.2(107)	125	8.8 (103)	103	1.9 (103)	110	6.4 (76)	120	1.7(100)	254	79.1(130)
6105	20	* *	16	* *	25	* *	13	* *	15	* *	72	* *
6106	1	* *	0	* *	0	* *	0	* *	0	* *	2	* *
6301	1,808	53.3 (90)	1,288	17.8 (60)	972	8.2 (62)	1,211	9.3 (65)	1,238	4.9 (59)	2,956	81.9(119)
											6,841	3.8 (44)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 56
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
6312	1,107	62.7 (63)	881	26.3 (18)	590	12.2 (19)	768	14.6 (40)	850	7.9 (37)	1,794	78.1(135)	4,203	3.5 (28)
6316	628	59.4 (73)	473	18.4 (54)	382	9.4 (45)	406	13.1 (44)	455	6.2 (50)	954	78.9(131)	2,338	2.7 (19)
6320	62	* *	58	* *	44	* *	44	* *	56	* *	228	92.1 (27)	385	0.5 (3)
6325	44	* *	41	* *	35	* *	33	* *	37	* *	135	84.4 (98)	254	5.1 (68)
6330	149	49.0 (99)	92	* *	73	* *	78	* *	86	* *	325	77.8(137)	638	9.2 (146)
6335	187	51.9 (92)	158	19.6 (47)	144	11.8 (21)	132	10.6 (56)	154	8.4 (31)	538	96.3 (5)	984	0.7 (6)
6340	50	* *	35	* *	23	* *	31	* *	33	* *	119	96.6 (2)	213	0.9 (9)
6345	4	* *	8	* *	4	* *	5	* *	8	* *	25	* *	45	* *
6350	76	* *	59	* *	78	* *	54	* *	57	* *	206	90.3 (52)	429	4.2 (51)
7304	67	* *	46	* *	56	* *	34	* *	38	* *	195	90.8 (41)	357	10.4 (159)
7306	960	50.9 (96)	583	17.7 (62)	570	3.0 (97)	507	12.2 (51)	503	6.6 (48)	1,954	83.7(103)	3,956	8.0 (130)
7308	56	* *	41	* *	43	* *	38	* *	27	* *	138	89.9 (57)	279	8.6 (137)
7310	23	* *	9	* *	13	* *	9	* *	9	* *	30	* *	75	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 57
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
7401	920	58.3 (76)	817	14.1 (80)	714	4.9 (87)	601	11.3 (55)	769	4.0 (67)
7601	244	65.6 (54)	232	21.1 (41)	168	13.1 (15)	196	17.9 (29)	227	4.4 (64)
7850	390	69.0 (35)	332	17.8 (60)	300	7.3 (70)	259	12.0 (52)	301	1.7 (100)
7852	41	* *	29	* *	22	* *	28	* *	29	* *
8101	436	44.7(103)	108	13.0 (85)	68	* *	136	8.8 (67)	100	2.0 (91)
8102	31	* *	16	* *	11	* *	16	* *	15	* *
8105	106	60.4 (68)	25	* *	22	* *	34	* *	23	* *
8106	62	* *	21	* *	17	* *	19	* *	19	* *
8150	14	* *	12	* *	10	* *	14	* *	12	* *
8151	1,058	68.0 (41)	871	31.7 (6)	769	15.6 (5)	708	18.1 (28)	817	11.1 (13)
8152	146	63.7 (57)	87	* *	64	* *	79	* *	83	* *
8153	13	* *	8	* *	4	* *	7	* *	6	* *
8154	121	75.2 (12)	93	* *	80	* *	85	* *	90	* *
									277	96.8 (1)
									519	1.3 (11)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 58
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)						Exclusive Breastfeeding (7)		≤2 Hours TV Viewing/Day (8)	Smoking in Household
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
8155	26	* *	19	* *	22	* *	20	* *	17	* *
8156	5	* *	6	* *	3	* *	3	* *	5	* *
8157	2	* *	4	* *	5	* *	1	* *	4	* *
8201	973	56.9 (81)	716	16.5 (69)	651	7.1 (74)	542	10.3 (57)	684	5.4 (55)
8202	1,202	64.1 (56)	885	29.0 (10)	777	10.4 (30)	741	5.3 (78)	848	3.2 (78)
8203	135	63.0 (59)	113	15.0 (74)	109	9.2 (49)	91	* *	109	3.7 (72)
8250	391	46.0(102)	284	13.0 (85)	234	11.1 (23)	209	9.6 (62)	225	4.4 (64)
8252	46	* *	15	* *	13	* *	20	* *	12	* *
8255	1,184	51.4 (94)	794	14.1 (80)	674	5.0 (85)	644	10.1 (58)	779	3.1 (81)
8257	148	56.8 (82)	120	25.8 (20)	131	13.7 (12)	100	3.0 (90)	113	1.8 (96)
8258	397	50.4 (97)	368	10.1 (99)	266	4.1 (91)	315	8.3 (70)	360	3.9 (69)
8260	313	71.9 (23)	230	35.2 (3)	224	23.7 (1)	192	7.3 (73)	223	3.6 (75)
8901	432	29.9(115)	333	9.3 (101)	254	2.0 (101)	309	4.5 (81)	310	2.3 (90)
									809	71.2(151)
										1,792 10.7 (162)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 59
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)	Smoking in Household
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
8902	359	40.7(110)	237	11.8 (92)	175	2.9 (98)	223	4.0 (87)	225	1.8 (96)
8903	478	23.6(117)	365	7.9 (105)	297	1.0 (105)	325	2.2 (94)	337	0.6(108)
8904	491	42.8(106)	379	11.6 (93)	286	4.5 (88)	356	4.5 (81)	359	1.9 (93)
8905	1,008	38.0(113)	756	10.4 (96)	658	3.5 (94)	659	2.7 (92)	696	1.7(100)
9004	1	* *	0	* *	0	* *	0	* *	0	* *
9010	6	* *	0	* *	0	* *	0	* *	0	* *
9012	15	* *	10	* *	9	* *	10	* *	10	* *
9013	48	* *	33	* *	25	* *	33	* *	32	* *
9015	19	* *	8	* *	6	* *	10	* *	8	* *
9016	69	* *	62	* *	39	* *	63	* *	62	* *
9017	27	* *	18	* *	11	* *	14	* *	17	* *
9018	50	* *	28	* *	22	* *	31	* *	27	* *
9019	16	* *	9	* *	8	* *	5	* *	7	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 60
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)									
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number	% (Rank)
9021	34	* *	34 * *	20 * *	24 * *	33 * *	78 * *	157	6.4 (91)
9040	0	* *	0 * *	0 * *	0 * *	0 * *	1 * *	1	* *
9041	9	* *	9 * *	7 * *	6 * *	8 * *	76 * *	103	9.7 (152)
9045	563	41.9(108)	437 10.3 (98)	454 3.5 (94)	389 3.1 (89)	412 2.4 (89)	1,314 69.3(156)	2,602	7.2 (117)
9050	475	38.5(112)	393 14.5 (77)	417 6.0 (79)	312 2.6 (93)	378 1.9 (93)	1,221 62.9(158)	2,432	8.6 (137)
9051	552	48.7(100)	399 22.1 (37)	358 8.4 (56)	323 5.0 (80)	370 3.5 (76)	1,462 76.5(140)	2,660	3.6 (31)
9053	938	43.8(105)	685 10.4 (96)	433 4.4 (89)	614 4.4 (85)	670 2.8 (83)	1,268 80.0(126)	3,091	21.3 (183)
9055	12	* *	0 * *	0 * *	0 * *	0 * *	1 * *	13	* *
9058	265	58.5 (75)	197 17.3 (65)	223 7.2 (71)	169 0.6 (96)	191 1.0(107)	750 93.5 (18)	1,345	6.0 (83)
9066	627	38.9(111)	560 5.4 (108)	435 2.1 (100)	472 1.7 (95)	555 1.4(103)	1,248 72.4(148)	2,699	9.2 (146)
9068	428	44.4(104)	365 8.8 (103)	350 3.1 (96)	296 4.4 (85)	360 2.8 (83)	881 75.0(141)	1,912	13.2 (174)
9069	699	57.4 (80)	603 9.5 (100)	450 4.2 (90)	512 3.9 (88)	594 1.3(104)	992 73.8(144)	2,589	9.2 (146)
9070	517	37.9(114)	464 6.0 (106)	450 2.0 (101)	339 7.1 (74)	439 2.5 (87)	1,085 59.4(160)	2,392	8.8 (143)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 61
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
9072	520	40.8(109)	404	9.2 (102)	332	2.4 (99)	358	2.8 (91)	395	1.3(104)
9074	375	55.5 (86)	331	12.1 (91)	220	5.0 (85)	276	6.2 (77)	323	1.9 (93)
9080	976	50.4 (97)	689	22.9 (31)	688	7.8 (67)	618	5.2 (79)	641	2.7 (86)
Unknown Clinics	140	60.0 (70)	84	* *	74	* *	78	* *	80	* *
Michigan	60,153	59.5	46,409	18.4	41,436	8.5	38,333	12.2	43,955	5.9
Nation (Prior Year)	1,689,615	63.2	908,604	25.1	1,069,942	16.9	363,039	10.7	466,279	6.1
									1,200,700	79.2
									4,203,554	9.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 62
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0101	114	4.4 (3)	9.6 (139)	594	4.9 (62)	2.5(108)	8.8 (33)	283	18.0(124)	11.7 (47)	390	12.8 (111)
0102	413	6.5 (27)	11.6 (164)	1,810	6.7(159)	4.1(178)	13.2(164)	826	18.2(130)	15.3(149)	1,101	13.5 (119)
0103	402	5.7 (13)	10.7 (158)	1,818	5.9(121)	2.0 (74)	13.9(174)	856	19.7(159)	18.1(172)	1,158	9.4 (69)
0104	128	7.8 (63)	10.2 (153)	606	5.9(121)	2.8(124)	11.4(102)	281	18.1(128)	12.8 (68)	361	6.4 (38)
0250	149	10.7 (140)	10.1 (149)	759	4.5 (44)	3.3(149)	12.1(131)	358	19.6(157)	17.0(168)	529	10.8 (82)
0251	110	7.3 (48)	6.4 (54)	603	3.6 (17)	0.8 (3)	13.4(168)	303	19.5(154)	19.5(177)	385	6.5 (39)
0252	113	6.2 (22)	9.7 (141)	679	4.0 (27)	1.6 (36)	13.8(171)	369	22.0(181)	18.2(173)	454	9.9 (74)
0253	153	8.5 (93)	11.8 (165)	786	2.7 (4)	9.2(204)	4.8 (1)	376	10.4 (4)	5.1 (2)	518	9.1 (67)
0401	535	9.2 (114)	12.3 (166)	2,441	6.0(126)	1.9 (67)	11.7(114)	1,143	16.4 (74)	15.2(144)	1,492	6.3 (33)
0402	372	5.1 (8)	8.1 (98)	1,928	4.4 (39)	2.1 (80)	12.8(155)	961	14.8 (39)	16.8(166)	1,249	11.0 (88)
0403	126	8.7 (99)	7.9 (92)	682	4.8 (55)	2.6(113)	15.2(189)	363	15.7 (54)	21.2(181)	445	2.9 (5)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 63
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature		Under- weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0404	131	8.4 (90)	9.2 (127)	716	2.9 (7)	2.4 (98)	10.6 (74)	377	15.6 (50)	15.1 (137)	444	10.8 (82)
0501	515	8.7 (99)	9.1 (121)	2,534	4.9 (62)	2.5 (108)	10.6 (74)	1,226	17.5 (113)	13.1 (76)	1,643	6.3 (33)
0502	0	* *	* *	13	* *	* *	* *	6	* *	* *	13	* *
0503	182	11.0 (143)	6.6 (62)	789	5.6 (100)	5.7 (192)	8.6 (24)	361	10.5 (5)	8.0 (9)	492	8.5 (60)
0504	516	8.3 (83)	10.1 (149)	2,675	5.8 (113)	1.7 (48)	12.2 (133)	1,301	19.7 (159)	13.5 (86)	1,785	5.2 (20)
0505	355	5.9 (15)	9.3 (133)	1,807	5.5 (93)	2.7 (120)	10.1 (60)	912	16.9 (91)	11.5 (44)	1,231	4.3 (14)
0506	161	6.8 (35)	6.8 (70)	981	4.9 (62)	4.0 (175)	11.6 (111)	544	17.3 (106)	12.9 (70)	689	5.4 (23)
0650	1,044	7.5 (55)	6.3 (51)	4,667	6.2 (137)	2.1 (80)	11.8 (120)	2,108	16.8 (90)	16.7 (165)	2,983	6.3 (33)
0651	653	6.7 (31)	6.0 (44)	3,003	6.0 (126)	2.3 (91)	11.8 (120)	1,430	17.1 (99)	14.3 (107)	1,962	10.7 (80)
0652	278	8.3 (83)	10.1 (149)	1,638	7.3 (179)	2.1 (80)	12.5 (146)	868	19.0 (146)	14.6 (116)	1,087	6.7 (41)
0653	417	9.6 (121)	8.6 (106)	2,225	5.8 (113)	2.0 (74)	12.3 (136)	1,134	16.7 (86)	15.2 (144)	1,493	10.9 (85)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 64
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0654	389	8.5 (93)	11.1 (161)	1,853	4.4 (39)	1.9 (67)	9.2 (37)	914	16.0 (61)	13.3 (79)	1,258	19.2 (151)
0655	375	6.4 (24)	6.7 (66)	1,728	7.3(179)	1.8 (55)	10.8 (78)	791	15.5 (49)	13.7 (92)	1,115	5.7 (27)
0657	64	* *	* *	401	5.7(104)	1.0 (7)	11.7(114)	228	19.3(152)	14.9(130)	294	19.4 (153)
0701	391	6.1 (19)	9.5 (138)	1,860	6.5(151)	1.8 (55)	8.0 (14)	895	12.1 (12)	9.6 (18)	1,037	7.8 (56)
0702	86	* *	* *	444	6.5(151)	2.7(120)	11.0 (86)	217	19.8(163)	15.7(155)	246	5.7 (27)
0703	216	5.1 (8)	5.1 (29)	1,059	5.7(104)	1.6 (36)	10.8 (78)	510	18.6(141)	14.5(112)	651	1.8 (1)
0704	72	* *	* *	428	6.3(140)	1.4 (19)	14.5(184)	246	19.5(154)	14.6(116)	268	3.7 (10)
0705	150	6.0 (17)	9.3 (133)	857	4.7 (50)	2.0 (74)	8.2 (16)	442	18.6(141)	7.9 (8)	487	3.9 (12)
0850	1,130	6.1 (19)	8.1 (98)	5,169	5.4 (88)	2.8(124)	11.5(109)	2,293	18.5(138)	14.8(127)	3,254	5.2 (20)
0851	835	6.6 (28)	8.5 (104)	4,283	4.4 (39)	2.5(108)	9.3 (39)	2,162	16.4 (74)	11.2 (41)	2,836	5.1 (19)
0901	1,722	8.3 (83)	9.2 (127)	8,060	6.3(140)	1.5 (29)	10.8 (78)	3,743	16.7 (86)	14.5(112)	5,360	11.3 (92)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 65
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0902	73	* *	* *	401	9.5(204)	2.7(120)	11.0 (86)	182	16.5 (76)	14.8(127)	276	21.7 (165)
1051	1,424	7.9 (68)	9.8 (143)	6,129	5.1 (72)	5.8(194)	11.0 (86)	2,916	16.6 (81)	12.7 (65)	3,733	10.9 (85)
1101	1,927	9.9 (124)	6.6 (62)	8,274	6.5(151)	2.0 (74)	10.7 (77)	3,780	16.6 (81)	13.6 (88)	5,377	15.2 (135)
1102	935	7.3 (48)	8.6 (106)	4,014	5.4 (88)	1.4 (19)	11.7(114)	1,810	17.1 (99)	14.4(110)	2,576	7.0 (43)
1106	109	3.7 (1)	7.3 (80)	517	4.3 (32)	0.2 (1)	10.1 (60)	239	17.2(102)	12.1 (54)	326	7.4 (50)
1250	1,022	6.8 (35)	9.0 (118)	5,196	6.4(144)	2.4 (98)	13.8(171)	2,566	17.9(122)	14.1(105)	3,351	5.4 (23)
1251	819	7.4 (53)	8.8 (112)	4,161	5.9(121)	1.8 (55)	10.4 (65)	2,027	18.3(133)	13.4 (82)	2,785	2.9 (5)
1252	114	7.9 (68)	8.8 (112)	881	5.4 (88)	1.6 (36)	19.8(200)	472	22.2(183)	26.3(188)	639	11.0 (88)
1253	1,344	8.1 (73)	8.2 (100)	6,349	7.3(179)	3.0(134)	12.5(146)	3,007	19.3(152)	15.1(137)	4,210	11.3 (92)
1350	2,378	8.0 (70)	7.8 (91)	11,090	6.6(156)	2.4 (98)	14.0(175)	5,136	18.4(136)	15.7(155)	7,280	14.4 (128)
1351	148	8.1 (73)	11.5 (162)	750	5.9(121)	1.6 (36)	14.5(184)	356	12.6 (16)	16.0(159)	518	10.0 (76)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 66
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1352	413	8.0 (70)	8.5 (104)	2,057	5.5 (93)	2.6(113)	13.2(164)	1,025	15.8 (56)	14.9(130)	1,383	22.6 (171)
1401	218	8.7 (99)	8.7 (109)	1,155	4.8 (55)	4.3(181)	12.5(146)	578	14.4 (33)	14.9(130)	761	3.8 (11)
1402	158	6.3 (23)	12.7 (168)	823	3.6 (17)	2.7(120)	14.1(178)	434	19.8(163)	13.4 (82)	573	8.9 (66)
1403	24	* *	* *	160	9.4(203)	1.9 (67)	12.5(146)	91	* *	* *	116	2.6 (3)
1404	14	* *	* *	93	* *	* *	* *	47	* *	* *	79	* *
1501	1,038	6.6 (28)	8.7 (109)	5,587	4.7 (50)	2.3 (91)	14.9(188)	2,853	17.4(108)	15.1(137)	3,886	23.2 (174)
1503	1,123	7.3 (48)	8.4 (102)	6,160	4.9 (62)	3.2(144)	12.2(133)	3,066	15.0 (42)	13.3 (79)	4,173	30.8 (184)
1504	2,292	7.2 (46)	6.8 (70)	11,572	4.2 (30)	3.2(144)	11.4(102)	5,761	16.3 (71)	13.9 (98)	7,810	19.3 (152)
1505	858	7.8 (63)	10.7 (158)	4,403	4.5 (44)	3.0(134)	10.4 (65)	2,289	13.8 (21)	9.9 (20)	2,881	22.3 (170)
1506	612	7.8 (63)	8.8 (112)	3,098	6.1(133)	3.4(155)	13.2(164)	1,512	19.7(159)	14.0(100)	2,079	21.2 (161)
1507	608	7.6 (56)	9.4 (136)	3,134	3.3 (10)	3.0(134)	8.7 (29)	1,536	17.4(108)	10.7 (32)	2,083	29.2 (181)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 67
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1508	102	4.9 (6)	4.9 (25)	655	2.7 (4)	2.4 (98)	18.9(198)	338	23.1(184)	21.3(182)	432	20.6 (157)
1509	597	8.9 (105)	7.7 (88)	3,343	3.9 (24)	2.8(124)	12.8(155)	1,766	18.5(138)	14.9(130)	2,305	18.1 (149)
1510	15	* *	* *	161	3.7 (20)	1.9 (67)	13.0(159)	77	* *	* *	112	12.5 (107)
1511	747	7.1 (43)	8.8 (112)	3,326	3.4 (14)	2.4 (98)	11.5(109)	1,522	18.2(130)	15.3(149)	2,106	22.2 (169)
1512	303	10.6 (139)	7.6 (85)	1,738	4.3 (32)	2.2 (86)	12.5(146)	902	18.0(124)	14.3(107)	1,181	26.7 (177)
1701	698	5.9 (15)	12.3 (166)	3,263	4.9 (62)	2.6(113)	13.0(159)	1,505	17.4(108)	14.6(116)	2,099	6.6 (40)
1712	27	* *	* *	171	7.6(188)	5.3(191)	12.9(158)	103	10.7 (6)	14.6(116)	130	14.6 (132)
1950	713	7.2 (46)	6.5 (58)	3,089	7.0(172)	1.9 (67)	11.2 (97)	1,366	18.5(138)	13.7 (92)	1,988	5.4 (23)
1952	131	5.3 (10)	8.4 (102)	738	4.9 (62)	2.6(113)	11.1 (92)	369	14.1 (27)	13.0 (73)	512	6.3 (33)
1953	269	6.7 (31)	6.7 (66)	1,535	5.8(113)	1.7 (48)	11.7(114)	774	14.7 (37)	13.8 (95)	1,040	4.8 (16)
1954	572	8.4 (90)	6.5 (58)	2,981	6.8(165)	2.1 (80)	12.3(136)	1,404	15.1 (44)	14.1(105)	1,847	5.8 (30)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 68
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1955	130	6.9 (37)	9.2 (127)	773	4.4 (39)	1.6 (36)	13.3(167)	410	13.9 (23)	15.1(137)	492	6.9 (42)
1957	533	7.7 (58)	9.2 (127)	2,418	6.0(126)	2.6(113)	12.3(136)	1,079	16.7 (86)	14.7(121)	1,575	4.5 (15)
1959	90	* *	* *	602	7.0(172)	2.8(124)	13.1(161)	303	12.5 (15)	15.2(144)	410	4.1 (13)
2150	658	8.2 (79)	9.1 (121)	3,243	6.7(159)	1.2 (9)	12.0(129)	1,563	16.1 (65)	15.2(144)	2,072	2.3 (2)
2151	261	7.7 (58)	9.2 (127)	1,194	4.6 (48)	1.7 (48)	10.5 (70)	543	19.0(146)	12.9 (70)	767	2.7 (4)
2251	439	8.0 (70)	7.7 (88)	2,031	5.8(113)	1.4 (19)	11.1 (92)	1,010	16.5 (76)	13.6 (88)	1,313	5.9 (31)
2252	220	7.3 (48)	5.9 (43)	1,043	5.7(104)	2.2 (86)	12.5(146)	506	21.1(175)	15.6(153)	696	5.7 (27)
2551	2,102	12.2 (152)	3.0 (3)	8,823	7.3(179)	3.1(138)	12.7(154)	3,762	16.9 (91)	13.4 (82)	5,681	21.9 (167)
2552	6	* *	* *	9	* *	* *	* *	1	* *	* *	2	* *
2554	103	8.7 (99)	3.9 (13)	638	5.8(113)	1.7 (48)	16.9(195)	334	19.5(154)	16.2(161)	417	14.4 (128)
2555	5,984	10.3 (131)	5.6 (36)	24,496	6.1(133)	3.2(144)	11.1 (92)	10,235	15.6 (50)	13.0 (73)	15,027	17.0 (145)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 69
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
3001	919	11.8 (150)	3.9 (13)	4,687	6.3(140)	2.3 (91)	15.6(191)	2,432	19.1(148)	15.6(153)	3,191 10.7 (80)
3002	242	11.6 (148)	5.0 (26)	1,318	6.2(137)	1.4 (19)	14.3(182)	707	20.9(173)	14.6(116)	908 17.8 (148)
3003	4	* *	* *	34	* *	* *	* *	19	* *	* *	28 * *
3004	126	4.8 (5)	1.6 (1)	900	5.7(104)	1.3 (13)	22.0(205)	536	21.8(180)	23.3(184)	683 7.9 (58)
3007	21	* *	* *	185	3.2 (9)	1.6 (36)	20.0(201)	107	20.6(171)	20.6(180)	152 13.8 (124)
3009	13	* *	* *	53	* *	* *	* *	29	* *	* *	44 * *
3010	2	* *	* *	8	* *	* *	* *	5	* *	* *	8 * *
3201	580	8.1 (73)	9.8 (143)	2,720	4.9 (62)	3.0(134)	12.3(136)	1,261	16.1 (65)	14.0(100)	1,624 10.8 (82)
3301	3,898	8.1 (73)	7.1 (73)	16,317	5.5 (93)	3.1(138)	12.3(136)	7,057	16.5 (76)	14.7(121)	10,364 11.5 (94)
3302	255	8.6 (96)	7.5 (83)	1,542	6.5(151)	1.8 (55)	14.1(178)	800	19.8(163)	17.5(171)	1,070 14.8 (133)
3304	127	7.1 (43)	10.2 (153)	714	3.9 (24)	3.4(155)	8.5 (21)	377	15.1 (44)	10.6 (30)	505 10.3 (78)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 70
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature		Under- weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
3305	149	5.4 (11)	10.1 (149)	724	4.1 (29)	3.6(164)	8.0 (14)	357	16.2 (67)	7.3 (5)	479	6.3 (33)
3306	267	8.2 (79)	12.7 (168)	1,295	5.2 (77)	3.2(144)	8.5 (21)	606	17.8(120)	8.7 (12)	851	7.4 (50)
3308	65	* *	* *	316	3.5 (16)	1.3 (13)	10.8 (78)	139	18.7(144)	10.1 (23)	207	7.7 (55)
3309	45	* *	* *	260	2.7 (4)	1.5 (29)	12.3(136)	137	21.2(177)	16.8(166)	183	4.9 (18)
3550	83	* *	* *	433	4.8 (55)	1.8 (55)	21.2(203)	231	21.2(177)	23.8(185)	270	13.3 (117)
3551	20	* *	* *	127	0.8 (1)	2.4 (98)	21.3(204)	71	* *	* *	84	* *
3552	46	* *	* *	233	6.4(144)	0.9 (4)	26.2(207)	122	17.2(102)	23.8(185)	135	23.7 (175)
3701	1,109	7.0 (39)	6.3 (51)	5,151	5.6(100)	3.3(149)	9.8 (47)	2,447	21.7(179)	11.9 (50)	3,575	8.6 (62)
3805	3,184	9.1 (112)	7.6 (85)	14,757	4.8 (55)	3.9(171)	9.8 (47)	6,847	17.3(106)	13.2 (77)	7,418	11.6 (99)
3950	1,905	7.6 (56)	9.0 (118)	8,903	4.6 (48)	4.0(175)	11.4(102)	4,112	17.9(122)	12.9 (70)	5,657	21.5 (164)
3952	5	* *	* *	67	* *	* *	* *	40	* *	* *	51	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 71
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
3953	4	* *	* *	78	* *	* *	* *	49	* *	* *	58 * *
3954	49	* *	* *	412	5.3 (83)	2.4 (98)	11.7(114)	242	16.5 (76)	12.4 (59)	297 14.5 (130)
4001	1,143	14.5 (165)	4.5 (19)	5,463	5.5 (93)	4.5(185)	12.4(143)	2,629	15.3 (47)	13.5 (86)	3,670 33.3 (187)
4002	1,041	10.3 (131)	5.6 (36)	5,060	5.3 (83)	3.9(171)	13.8(171)	2,421	17.1 (99)	15.8(157)	3,423 25.2 (176)
4103	89	* *	* *	585	4.4 (39)	4.4(183)	9.9 (52)	321	14.6 (35)	10.3 (25)	411 21.2 (161)
4106	1,612	10.5 (137)	5.7 (39)	7,559	6.4(144)	1.5 (29)	11.9(123)	3,674	16.0 (61)	14.7(121)	4,991 16.5 (144)
4107	3,132	9.4 (116)	7.9 (92)	14,231	6.8(165)	1.6 (36)	11.1 (92)	6,436	16.6 (81)	12.6 (62)	9,166 14.5 (130)
4109	1,142	8.3 (83)	9.4 (136)	6,102	4.8 (55)	1.9 (67)	8.3 (17)	3,058	15.6 (50)	10.9 (37)	4,106 7.5 (54)
4110	1,096	8.3 (83)	6.8 (70)	6,408	5.8(113)	1.2 (9)	14.1(178)	3,224	18.3(133)	17.3(170)	4,382 11.5 (94)
4112	568	3.7 (1)	9.7 (141)	3,256	6.4(144)	3.5(161)	19.2(199)	1,765	19.7(159)	21.8(183)	2,650 12.1 (103)
4113	672	4.9 (6)	8.8 (112)	3,404	3.3 (10)	1.7 (48)	14.0(175)	1,680	18.6(141)	19.0(176)	2,269 7.4 (50)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 72
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4120	2,015	8.3 (83)	8.3 (101)	10,120	7.0(172)	1.5 (29)	12.4(143)	4,857	16.9 (91)	13.0 (73)	6,717	13.5 (119)
4160	1,777	10.5 (137)	7.3 (80)	8,596	6.7(159)	1.6 (36)	11.4(102)	3,970	16.0 (61)	12.4 (59)	5,437	12.6 (109)
4351	540	7.8 (63)	10.7 (158)	2,920	5.7(104)	1.8 (55)	11.1 (92)	1,483	19.1(148)	14.0(100)	1,987	12.5 (107)
4352	168	10.1 (127)	7.1 (73)	915	6.7(159)	1.6 (36)	12.3(136)	487	20.7(172)	13.6 (88)	653	15.2 (135)
4353	433	8.5 (93)	9.0 (118)	2,432	5.7(104)	2.8(124)	11.7(114)	1,202	20.5(169)	14.0(100)	1,713	12.4 (106)
4354	794	5.7 (13)	9.9(147)	3,627	4.8 (55)	1.8 (55)	9.4 (43)	1,671	18.1(128)	11.5 (44)	2,353	9.4 (69)
4355	297	8.1 (73)	9.1 (121)	1,551	5.2 (77)	1.5 (29)	13.1(161)	767	21.1(175)	17.1(169)	1,049	9.6 (73)
4356	541	8.1 (73)	6.1 (47)	2,641	6.7(159)	1.5 (29)	11.9(123)	1,290	20.9(173)	15.1(137)	1,777	12.9 (112)
4357	3	* *	* *	8	* *	* *	* *	3	* *	* *	4	* *
4358	168	8.9(105)	5.4 (34)	1,063	7.1(175)	0.7 (2)	21.0(202)	603	26.7(188)	28.4(189)	770	12.6 (109)
4359	519	5.4 (11)	10.4 (156)	2,750	6.2(137)	1.7 (48)	15.6(191)	1,470	19.1(148)	15.3(149)	1,875	12.2 (105)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 73
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4360	165	6.7 (31)	9.1 (121)	849	3.8 (22)	1.4 (19)	8.7 (29)	419	16.2 (67)	10.5 (27)	575	13.2 (116)
4361	221	9.0 (108)	5.0 (26)	1,052	6.4(144)	0.9 (4)	11.3(100)	486	24.1(187)	16.5(163)	709	13.5 (119)
4362	313	9.6 (121)	8.6 (106)	1,562	7.1(175)	1.3 (13)	8.6 (24)	763	14.5 (34)	11.9 (50)	1,058	13.3 (117)
4363	869	7.4 (53)	9.2 (127)	4,188	5.1 (72)	1.3 (13)	9.7 (46)	1,986	16.3 (71)	13.8 (95)	2,833	15.3 (137)
4401	1,331	9.0 (108)	7.9 (92)	5,943	7.6(188)	2.8(124)	11.3(100)	2,696	17.2(102)	14.7(121)	3,792	9.9 (74)
4501	932	6.9 (37)	7.1 (73)	4,742	6.0(126)	2.0 (74)	8.4 (19)	2,312	14.7 (37)	10.5 (27)	3,301	7.2 (47)
4502	1	* *	* *	4	* *	* *	* *	2	* *	* *	3	* *
4591	0	* *	* *	3	* *	* *	* *	2	* *	* *	3	* *
4601	1,540	8.2 (79)	6.3 (51)	6,650	5.1 (72)	2.9(131)	12.1(131)	2,872	18.9(145)	15.5(152)	3,552	7.2 (47)
4701	1,350	6.6 (28)	10.2 (153)	5,667	5.2 (77)	1.8 (55)	10.1 (60)	2,510	17.4(108)	15.1(137)	3,706	5.3 (22)
5050	1,747	11.1 (144)	5.8 (42)	6,658	5.7(104)	2.4 (98)	11.9(123)	2,598	14.2 (30)	12.5 (61)	4,165	16.3 (142)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 74
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature		Under-weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
5051	2,927	10.2 (128)	6.2 (49)	11,073	6.6(156)	2.4 (98)	11.8(120)	4,332	13.9 (23)	10.8 (34)	6,950	17.2 (147)
5052	2,708	10.0 (125)	5.0 (26)	10,157	5.6(100)	1.7 (48)	12.2(133)	3,992	17.0 (95)	14.7(121)	6,380	15.1 (134)
5101	2,320	7.7 (58)	6.4 (54)	9,511	5.3 (83)	1.8 (55)	10.5 (70)	3,997	17.4(108)	13.7 (92)	5,882	10.5 (79)
5202	626	7.0 (39)	7.2 (78)	2,626	5.7(104)	2.0 (74)	13.1(161)	1,185	20.3(168)	16.5(163)	1,485	9.4 (69)
5203	150	6.0 (17)	9.3(133)	794	4.8 (55)	1.6 (36)	12.8(155)	402	16.2 (67)	13.2 (77)	496	10.1 (77)
5204	130	7.7 (58)	11.5(162)	563	4.3 (32)	1.4 (19)	10.8 (78)	267	19.1(148)	12.7 (65)	314	11.1 (90)
5601	1,252	8.9(105)	8.8(112)	5,621	7.2(178)	2.2 (86)	14.5(184)	2,610	15.8 (56)	14.4(110)	3,368	7.4 (50)
5801	2,036	9.4(116)	7.7 (88)	8,604	7.7(192)	1.5 (29)	11.6(111)	3,693	17.7(119)	14.7(121)	5,396	8.7 (63)
5803	143	7.7 (58)	9.8(143)	848	5.3 (83)	2.4 (98)	10.8 (78)	400	13.8 (21)	12.3 (58)	579	9.2 (68)
5804	46	* *	* *	407	3.4 (14)	3.7(166)	9.3 (39)	216	11.1 (7)	11.1 (39)	307	7.8 (56)
6101	65	* *	* *	397	6.0(126)	3.8(169)	8.6 (24)	219	17.8(120)	9.6 (18)	281	11.7(100)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 75
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature Under-weight Obese				≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
6102	193	7.3 (48)	6.7 (66)	1,192	5.4 (88)	3.1(138)	10.5 (70)	632	15.7 (54)	12.7 (65)	780	12.9 (112)
6103	3,760	9.0 (108)	6.5 (58)	17,032	4.7 (50)	3.4(155)	10.3 (64)	8,101	16.5 (76)	12.2 (57)	11,096	20.4 (156)
6104	466	10.3 (131)	5.2 (30)	1,377	4.2 (30)	5.7(192)	9.9 (52)	593	17.2(102)	15.0(136)	755	15.9 (140)
6105	53	* *	* *	383	3.7 (20)	2.3 (91)	6.8 (4)	214	15.9 (59)	8.4 (11)	280	13.6 (122)
6106	22	* *	* *	79	* *	* *	* *	40	* *	* *	48	* *
6301	4,892	9.6 (121)	6.4 (54)	19,955	6.1(133)	1.8 (55)	10.0 (57)	8,376	16.0 (61)	14.5(112)	12,692	12.9 (112)
6312	2,828	10.3 (131)	6.5 (58)	11,555	6.9(170)	2.2 (86)	9.3 (39)	4,775	16.3 (71)	12.1 (54)	7,350	17.1 (146)
6316	1,612	8.7 (99)	7.9 (92)	6,386	5.1 (72)	1.8 (55)	8.7 (29)	2,492	14.6 (35)	12.1 (54)	4,017	9.5 (72)
6320	208	8.2 (79)	9.1 (121)	1,230	4.3 (32)	4.1(178)	9.8 (47)	651	13.1 (19)	11.8 (49)	797	7.2 (47)
6325	134	9.0 (108)	6.0 (44)	738	6.0(126)	3.5(161)	9.9 (52)	354	22.0(181)	13.3 (79)	487	6.0 (32)
6330	405	15.1 (166)	5.7 (39)	1,770	6.8(165)	2.9(131)	11.0 (86)	818	14.2 (30)	11.5 (44)	1,104	18.9 (150)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 76
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
6335	529	7.0 (39)	9.1 (121)	2,809	3.6 (17)	3.4(155)	8.5 (21)	1,376	14.8 (39)	9.3 (15)	1,791	8.8 (64)
6340	127	9.4 (116)	4.7 (20)	691	3.9 (24)	1.6 (36)	7.8 (11)	378	11.4 (8)	10.6 (30)	442	8.8 (64)
6345	26	* *	* *	169	7.7(192)	1.2 (9)	16.0(194)	80	* *	* *	104	11.5 (94)
6350	254	9.1 (112)	7.1 (73)	1,246	5.4 (88)	3.4(155)	11.9(123)	529	14.0 (25)	15.1(137)	767	14.3 (126)
7304	202	6.4 (24)	9.9(147)	1,115	4.3 (32)	2.8(124)	15.2(189)	571	23.6(185)	20.5(179)	784	2.9 (5)
7306	2,555	10.3 (131)	7.4 (82)	11,422	5.2 (77)	2.5(108)	12.0(129)	5,335	17.0 (95)	14.9(130)	7,799	13.0 (115)
7308	183	7.1 (43)	9.8(143)	933	4.7 (50)	2.3 (91)	11.6(111)	451	16.9 (91)	14.9(130)	636	8.5 (60)
7310	63	* *	* *	206	5.8(113)	3.9(171)	11.2 (97)	84	* *	* *	127	5.5 (26)
7401	2,397	8.4 (90)	6.7 (66)	10,841	6.0(126)	2.1 (80)	9.0 (34)	4,960	15.4 (48)	11.3 (43)	7,101	8.3 (59)
7402	49	* *	* *	324	8.0(198)	2.5(108)	11.4(102)	143	11.9 (9)	20.3(178)	244	11.1 (90)
7403	21	* *	* *	110	6.4(144)	3.6(164)	9.1 (36)	52	* *	* *	77	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 77
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
7404	16	* *	* *	100	5.0 (71)	4.0(175)	10.0 (57)	50	* *	* *	80	* *
7601	704	6.1 (19)	5.3 (31)	3,316	4.0 (27)	3.5(161)	7.1 (7)	1,507	12.2 (13)	9.4 (16)	2,261	4.8 (16)
7850	1,021	8.3 (83)	5.6 (36)	4,763	4.3 (32)	5.1(189)	11.4(102)	2,366	17.5(113)	15.8(157)	3,218	3.0 (8)
7851	47	* *	* *	258	6.6(156)	3.1(138)	12.4(143)	137	18.2(130)	18.2(173)	184	7.1 (46)
7852	105	6.7 (31)	10.5 (157)	459	3.3 (10)	5.0(186)	9.8 (47)	216	17.6(118)	13.4 (82)	309	3.2 (9)
8101	434	22.1 (169)	4.4 (18)	713	7.4(184)	11.1(206)	8.7 (29)	205	12.2 (13)	11.2 (41)	307	32.6 (186)
8102	31	* *	* *	134	8.2(199)	6.0(195)	14.2(181)	75	* *	* *	99	* *
8105	103	8.7 (99)	8.7 (109)	216	10.6(206)	1.4 (19)	12.5(146)	67	* *	* *	94	* *
8106	62	* *	* *	164	4.9 (62)	3.7(166)	14.0(175)	86	* *	* *	109	30.3 (183)
8150	47	* *	* *	341	3.8 (22)	2.6(113)	7.0 (5)	196	15.8 (56)	8.7 (12)	228	15.4 (138)
8151	2,884	9.3 (115)	6.6 (62)	12,307	5.5 (93)	2.2 (86)	11.0 (86)	5,293	16.6 (81)	13.6 (88)	7,844	22.1 (168)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 78
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8152	437	11.9 (151)	6.4 (54)	1,315	5.9(121)	5.0(186)	10.9 (85)	394	17.0 (95)	16.0(159)	720	39.9 (190)
8153	29	* *	* *	151	7.9(196)	1.3 (13)	11.9(123)	87	* *	* *	104	11.5 (94)
8154	281	6.4 (24)	9.6 (139)	1,416	3.0 (8)	3.9(171)	6.5 (2)	689	15.1 (44)	8.3 (10)	870	7.0 (43)
8155	57	* *	* *	397	1.3 (2)	2.3 (91)	8.6 (24)	223	17.5(113)	7.6 (6)	286	19.9 (154)
8156	17	* *	* *	140	7.1(175)	1.4 (19)	8.6 (24)	79	* *	* *	101	10.9 (85)
8157	15	* *	* *	116	4.3 (32)	0.9 (4)	9.5 (45)	78	* *	* *	75	* *
8201	2,672	8.6 (96)	7.5 (83)	8,519	6.8(165)	1.8 (55)	10.4 (65)	2,322	18.3(133)	11.9 (50)	4,448	7.0 (43)
8202	2,750	7.8 (63)	7.2 (78)	13,599	4.5 (44)	1.9 (67)	11.0 (86)	6,547	16.7 (86)	12.6 (62)	8,732	22.6 (171)
8203	423	10.4 (136)	7.6 (85)	1,456	7.6(188)	1.2 (9)	10.0 (57)	487	20.5(169)	9.4 (16)	769	16.1 (141)
8250	1,117	10.8 (141)	6.0 (44)	3,143	5.7(104)	3.3(149)	7.6 (10)	956	12.8 (18)	10.0 (22)	1,511	20.9 (159)
8252	45	* *	* *	127	11.0(207)	7.9(203)	7.1 (7)	65	* *	* *	75	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 79
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature		Under- weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8255	3,094	9.4 (116)	7.9 (92)	10,063	7.9(196)	1.0 (7)	11.2 (97)	2,957	20.2(167)	14.0(100)	5,313	13.6 (122)
8257	437	10.8 (141)	5.5 (35)	2,349	5.5 (93)	4.4(183)	10.4 (65)	1,125	12.6 (16)	12.8 (68)	1,416	14.3 (126)
8258	1,271	9.4 (116)	6.1 (47)	5,875	6.9(170)	1.4 (19)	12.5(146)	2,284	18.4(136)	13.9 (98)	3,548	11.8 (101)
8260	636	10.2 (128)	3.1 (5)	2,765	5.2 (77)	3.4(155)	7.4 (9)	1,290	14.1 (27)	9.2 (14)	1,801	20.8 (158)
8901	1,098	14.0 (161)	3.6 (10)	4,768	6.1(133)	5.1(189)	10.4 (65)	2,140	13.7 (20)	11.1 (39)	2,913	38.8 (189)
8902	790	11.3 (146)	3.0 (3)	3,132	5.2 (77)	6.6(199)	9.9 (52)	1,367	11.9 (9)	10.8 (34)	1,938	32.1 (185)
8903	1,290	13.7 (160)	3.7 (11)	5,630	6.3(140)	5.0(186)	11.4(102)	2,544	15.9 (59)	12.0 (53)	3,579	41.6 (192)
8904	1,131	12.6 (156)	5.7 (39)	4,681	4.5 (44)	6.5(198)	9.3 (39)	2,081	14.2 (30)	10.7 (32)	3,005	37.7 (188)
8905	2,936	8.6 (96)	6.6 (62)	13,871	5.1 (72)	6.6(199)	13.5(170)	6,683	16.6 (81)	14.5(112)	8,449	28.8 (179)
9004	16	* *	* *	11	* *	* *	* *	0	* *	* *	0	* *
9010	6	* *	* *	3	* *	* *	* *	0	* *	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 80
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)											
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number % (Rank)
9012	32	* *	* *	189	1.6 (3)	2.1 (80)	7.9 (12)	99	* *	* *	33 * *
9013	106	17.9 (168)	4.7 (20)	349	4.9 (62)	3.7 (166)	8.3 (17)	143	19.6 (157)	7.7 (7)	116 40.5 (191)
9015	32	* *	* *	130	7.7 (192)	3.8 (169)	18.5 (197)	70	* *	* *	48 * *
9016	137	16.1 (167)	2.2 (2)	521	7.5 (187)	2.3 (91)	10.2 (63)	220	20.0 (166)	10.5 (27)	271 46.1 (193)
9017	56	* *	* *	231	10.0 (205)	1.3 (13)	17.7 (196)	112	27.7 (189)	14.3 (107)	41 * *
9018	128	10.2 (128)	3.1 (5)	410	9.3 (202)	6.8 (201)	9.8 (47)	165	17.0 (95)	16.4 (162)	20 * *
9019	35	* *	* *	162	7.4 (184)	4.3 (181)	10.5 (70)	83	* *	* *	14 * *
9021	82	* *	* *	314	7.3 (179)	6.1 (197)	14.6 (187)	138	23.9 (186)	13.8 (95)	34 * *
9040	6	* *	* *	20	* *	* *	* *	12	* *	* *	11 * *
9041	171	14.0 (161)	4.7 (20)	1,303	5.8 (113)	4.1 (178)	11.9 (123)	691	17.5 (113)	10.3 (25)	389 29.6 (182)
9045	1,413	11.2 (145)	7.1 (73)	6,769	6.4 (144)	1.6 (36)	14.4 (183)	3,308	17.5 (113)	14.8 (127)	3,559 22.8 (173)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 81
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature			Under-weight	Obese	≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9050	1,746	11.6 (148)	6.2 (49)	8,160	6.8(165)	3.3(149)	13.4(168)	3,753	14.8 (39)	15.2(144)	5,125	15.5 (139)
9051	1,609	7.0 (39)	7.9 (92)	8,679	5.3 (83)	1.4 (19)	15.7(193)	4,133	18.0(124)	18.5(175)	6,009	13.8 (124)
9053	2,406	14.4 (164)	4.1 (15)	9,157	6.7(159)	3.2(144)	9.2 (37)	3,652	16.2 (67)	11.7 (47)	5,772	21.7 (165)
9055	245	11.4 (147)	5.3 (31)	228	9.2(201)	14.9(207)	7.0 (5)	1	*	*	*	*
9058	659	10.0 (125)	5.3 (31)	3,672	3.3 (10)	3.3(149)	9.0 (34)	1,923	11.9 (9)	9.9 (20)	2,495	21.0 (160)
9066	1,930	13.1 (158)	4.2 (16)	8,291	5.5 (93)	6.0(195)	7.9 (12)	3,407	8.3 (3)	6.4 (4)	5,104	11.5 (94)
9068	1,395	12.4 (155)	3.4 (8)	5,709	5.6(100)	10.3(205)	6.6 (3)	2,406	5.7 (1)	4.2 (1)	3,673	28.6 (178)
9069	2,131	12.3 (154)	4.8 (24)	7,852	6.5(151)	3.1(138)	9.4 (43)	2,814	15.0 (42)	12.6 (62)	4,486	20.2 (155)
9070	1,785	12.8 (157)	3.1 (5)	7,353	7.6(188)	7.7(202)	10.6 (74)	3,100	7.2 (2)	5.9 (3)	4,474	21.2 (161)
9072	1,540	13.2 (159)	3.8 (12)	6,053	7.4(184)	2.6(113)	10.8 (78)	2,461	15.6 (50)	10.1 (23)	3,801	12.1 (103)
9074	1,259	14.2 (163)	3.5 (9)	5,327	7.8(195)	3.1(138)	9.9 (52)	2,274	14.1 (27)	10.9 (37)	3,340	16.3 (142)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 82
Run Date: 3 / 12 / 12

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9080	2,529	12.2 (152)	4.3 (17)	11,608	4.7 (50)	2.9(131)	8.4 (19)	5,509	14.0 (25)	10.8 (34)	7,530	29.1 (180)
9151	0	* *	* *	1	* *	* *	* *	1	* *	* *	2	* *
Unknown Clinics	169	4.7 (4)	4.7 (20)	719	8.6(200)	3.3(149)	22.3(206)	356	18.0(124)	25.8(187)	554	11.9 (102)
Michigan	164,970	9.3	6.9	741,082	5.8	2.8	11.3	336,514	16.5	13.4	470,756	15.2

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 83
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
0101	140	67.1 (48)	129	27.9 (16)	117	12.0 (35)	112	22.3 (23)	119	9.2 (32)	290	86.6 (98)	308	8.1 (118)
0102	514	64.8 (70)	394	13.7 (129)	341	4.7 (143)	364	14.6 (80)	363	3.3(126)	828	85.4(109)	1,004	8.5 (130)
0103	489	60.5 (94)	402	16.9 (109)	346	6.4 (121)	358	19.0 (41)	356	7.3 (64)	891	83.3(128)	1,000	11.3 (164)
0104	161	67.1 (48)	146	9.6 (156)	121	3.3 (154)	132	10.6 (108)	140	2.1(147)	275	78.2(157)	322	19.9 (191)
0250	170	65.3 (65)	168	20.8 (65)	160	12.5 (31)	143	18.2 (47)	156	10.9 (17)	358	88.0 (72)	394	8.4 (126)
0251	132	51.5(133)	126	11.9 (148)	106	6.6 (119)	109	15.6 (71)	109	6.4 (72)	294	88.4 (66)	300	7.7 (108)
0252	138	73.2 (20)	115	32.2 (9)	132	16.7 (9)	107	38.3 (2)	113	16.8 (3)	394	86.5(101)	351	10.0 (147)
0253	183	56.8(111)	172	14.5 (123)	147	3.4 (152)	164	15.2 (72)	168	8.9 (39)	394	87.1 (85)	408	12.3 (170)
0401	615	48.5(144)	542	18.8 (88)	492	9.1 (86)	517	19.3 (35)	528	8.9 (39)	1,153	92.7 (15)	1,266	7.6 (102)
0402	454	65.9 (57)	392	20.7 (69)	347	9.5 (77)	350	17.7 (53)	346	4.3(105)	972	87.6 (79)	1,017	6.9 (90)
0403	148	62.2 (91)	124	12.9 (138)	122	5.7 (131)	122	12.3 (96)	123	2.4(137)	377	84.1(122)	339	8.8 (133)
0404	165	49.7(141)	129	20.2 (72)	118	9.3 (81)	130	14.6 (80)	116	3.4(124)	374	86.4(104)	364	20.1 (192)
0501	655	71.1 (28)	550	25.6 (28)	474	11.6 (41)	503	28.4 (7)	511	11.5 (12)	1,259	90.2 (39)	1,371	12.2 (167)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 84
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
0502	2	* *	3	* *	4	* *	2	* *	3	* *
0503	222	80.2 (5)	187	32.6 (8)	152	14.5 (19)	177	27.1 (9)	177	14.7 (4)
0504	656	77.9 (9)	546	31.0 (11)	442	16.7 (9)	486	31.7 (4)	468	14.5 (5)
0505	441	77.6 (10)	379	29.8 (12)	322	16.1 (11)	365	29.9 (5)	363	9.4 (29)
0506	193	78.2 (8)	180	20.6 (70)	179	11.7 (40)	151	26.5 (11)	160	8.1 (48)
0650	1,252	69.2 (39)	1,108	21.3 (57)	973	10.1 (62)	983	19.2 (36)	1,033	8.0 (50)
0651	750	57.2(109)	650	13.5 (132)	605	6.0 (127)	619	14.4 (84)	629	3.5(123)
0652	339	61.7 (92)	303	21.1 (61)	279	7.9 (104)	271	15.1 (73)	276	6.5 (71)
0653	490	64.3 (74)	457	23.9 (35)	428	9.6 (75)	389	23.4 (21)	418	11.0 (14)
0654	446	66.1 (56)	395	17.5 (104)	379	10.0 (68)	370	16.8 (62)	380	7.6 (59)
0655	443	71.3 (25)	415	18.8 (88)	355	9.6 (75)	364	17.6 (54)	389	7.7 (55)
0657	79	* *	68	* *	65	* *	64	* *	65	* *
0701	493	81.3 (3)	429	39.4 (3)	355	19.2 (6)	400	26.5 (11)	414	13.3 (8)
	931	97.0 (1)	982	1.0 (5)						

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 85
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household		
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
0702	102	57.8(108)	98	* *	85	* *	84	* *	92	* *
0703	273	63.7 (79)	233	21.0 (63)	198	10.1 (62)	213	21.6 (24)	227	6.6 (68)
0704	87	* *	69	* *	65	* *	64	* *	63	* *
0705	193	69.9 (33)	179	25.7 (27)	143	9.8 (70)	163	16.6 (63)	175	9.7 (27)
0850	1,390	71.2 (27)	1,265	21.0 (63)	1,059	9.9 (69)	1,168	18.2 (47)	1,213	7.7 (55)
0851	1,033	64.6 (73)	962	18.7 (91)	864	8.0 (100)	872	19.7 (30)	933	7.6 (59)
0901	2,015	57.2(109)	1,783	13.7 (129)	1,577	5.6 (136)	1,712	13.6 (87)	1,656	3.0(128)
0902	92	* *	82	* *	73	* *	72	* *	69	* *
1051	1,704	81.3 (3)	1,311	34.0 (5)	1,161	17.3 (7)	1,149	28.1 (8)	1,240	11.0 (14)
1101	2,266	46.3(150)	1,807	13.0 (137)	1,680	7.0 (112)	1,637	9.6 (121)	1,718	4.4(104)
1102	1,139	52.0(129)	956	17.9 (99)	859	10.7 (52)	875	13.8 (86)	911	6.9 (66)
1106	143	72.0 (22)	120	26.7 (20)	106	13.2 (24)	107	13.1 (91)	112	4.5(102)
1250	1,240	58.8(105)	1,093	19.9 (77)	935	9.5 (77)	1,068	16.9 (61)	1,059	9.3 (31)
	2,733	84.4(120)	2,854	7.8 (112)						

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 86
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
1251	1,029	68.4 (41)	864	22.3 (51)	739	10.6 (53)	823	20.8 (28)	820	10.7 (18)	2,157	89.8 (44)	2,355	7.7 (108)
1252	57	* *	95	* *	103	3.9 (149)	62	* *	84	* *	519	80.3(144)	569	5.8 (67)
1253	1,721	63.6 (81)	1,443	19.2 (86)	1,245	8.1 (97)	1,395	15.0 (74)	1,356	9.6 (28)	3,213	87.1 (85)	3,549	5.9 (72)
1350	2,838	53.6(120)	2,334	12.9 (138)	1,976	5.0 (141)	2,268	9.7 (119)	2,248	4.0(114)	5,458	89.5 (49)	6,197	5.3 (58)
1351	168	69.6 (35)	149	18.8 (88)	148	10.8 (50)	148	9.5 (122)	147	4.8 (99)	397	96.5 (4)	437	3.2 (20)
1352	492	52.8(125)	415	9.6 (156)	393	4.3 (146)	400	6.3 (140)	410	2.4(137)	1,112	88.0 (72)	1,135	10.4 (155)
1401	248	69.4 (37)	227	22.9 (44)	210	8.1 (97)	211	20.9 (26)	214	8.9 (39)	583	88.7 (63)	583	4.8 (40)
1402	184	83.7 (2)	148	33.1 (6)	145	22.1 (4)	133	24.8 (16)	130	10.0 (23)	434	86.9 (93)	432	6.5 (81)
1403	29	* *	25	* *	27	* *	22	* *	24	* *	93	* *	84	* *
1404	19	* *	20	* *	13	* *	18	* *	20	* *	49	* *	60	* *
1501	1,283	68.2 (44)	1,108	26.3 (22)	1,044	11.5 (42)	1,037	19.7 (30)	1,011	7.8 (54)	3,013	85.0(114)	3,088	6.5 (81)
1503	1,463	65.4 (61)	1,262	23.4 (40)	1,178	13.1 (25)	1,134	17.0 (59)	1,144	9.4 (29)	3,271	84.7(118)	3,477	7.5 (100)
1504	2,916	70.3 (32)	2,288	24.6 (33)	2,266	11.4 (43)	1,989	18.9 (42)	2,047	7.7 (55)	6,045	85.2(111)	6,714	3.2 (20)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 87
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)											
Clinic	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household		
	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)	
1505	1,051	75.9 (15)	848 32.9 (7)	806 14.9 (15)	775 28.9 (6)	799 14.3 (6)	2,416	91.3 (26)	2,477	2.1 (11)	
1506	745	71.0 (29)	645 21.7 (55)	611 10.5 (55)	578 18.2 (47)	585 8.9 (39)	1,606	91.2 (27)	1,705	6.0 (73)	
1507	773	70.5 (31)	646 25.9 (26)	621 11.4 (43)	590 25.1 (14)	622 10.5 (20)	1,622	91.1 (29)	1,778	4.8 (40)	
1508	151	76.2 (14)	140 26.4 (21)	137 14.6 (17)	108 9.3 (125)	119 12.6 (9)	362	92.5 (16)	433	0.0 (1)	
1509	725	65.5 (58)	654 23.1 (43)	626 10.2 (58)	578 14.5 (83)	599 6.2 (74)	1,833	83.7(124)	1,825	6.1 (75)	
1510	11	* *	18 * *	25 * *	9 * *	18 * *	79	* *	111	12.6 (171)	
1511	875	70.7 (30)	761 18.4 (94)	632 9.7 (71)	744 17.6 (54)	743 5.4 (88)	1,618	88.4 (66)	1,880	4.9 (46)	
1512	416	62.5 (90)	375 21.3 (57)	326 11.0 (48)	362 18.2 (47)	367 9.8 (26)	952	92.1 (21)	976	4.1 (29)	
1701	847	65.4 (61)	705 22.8 (45)	575 11.8 (38)	647 25.0 (15)	631 10.6 (19)	1,527	87.6 (79)	1,676	6.6 (84)	
1712	10	* *	17 * *	19 * *	12 * *	15 * *	89	* *	86	* *	
1950	831	66.5 (54)	720 18.5 (92)	589 8.3 (93)	671 16.1 (66)	695 6.6 (68)	1,462	88.0 (72)	1,780	9.3 (139)	
1952	172	69.8 (34)	159 22.0 (53)	139 10.8 (50)	139 23.7 (19)	146 8.9 (39)	391	89.5 (49)	426	8.2 (119)	
1953	343	65.0 (69)	318 19.8 (79)	269 6.3 (122)	287 13.6 (87)	300 6.0 (77)	827	86.9 (93)	837	6.9 (90)	

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 88
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
1954	735	65.2 (67)	633	21.2 (60)	549	10.6 (53)	601	19.6 (32)	605	8.9 (39)	1,517	91.0 (31)	1,633	7.0 (93)
1955	160	69.4 (37)	129	19.4 (82)	124	11.3 (45)	121	16.5 (64)	120	5.8 (80)	436	87.6 (79)	415	8.0 (116)
1957	695	73.7 (18)	606	22.3 (51)	487	12.5 (31)	579	20.9 (26)	580	6.7 (67)	1,149	86.5(101)	1,392	4.0 (27)
1959	71	* *	102	22.5 (49)	101	15.8 (13)	72	* *	94	* *	318	87.4 (82)	306	4.9 (46)
2150	798	63.2 (85)	704	18.2 (96)	628	8.9 (90)	624	14.6 (80)	622	6.1 (76)	1,595	92.9 (14)	1,677	6.0 (73)
2151	316	60.1 (99)	280	16.4 (111)	242	9.1 (86)	271	15.9 (68)	272	4.0(114)	540	89.3 (53)	659	7.3 (97)
2251	516	63.8 (78)	442	23.5 (39)	385	11.2 (47)	405	23.7 (19)	423	8.0 (50)	985	89.0 (58)	1,082	4.6 (36)
2252	249	59.0(104)	223	23.8 (37)	199	10.1 (62)	203	27.1 (9)	213	10.3 (21)	502	91.2 (27)	527	6.6 (84)
2551	1,602	36.8(165)	1,866	7.2 (163)	1,579	3.1 (158)	1,620	6.9 (135)	1,731	1.6(157)	3,712	75.9(162)	4,871	8.8 (133)
2552	7	* *	1	* *	0	* *	1	* *	1	* *	1	* *	7	* *
2554	141	46.1(151)	131	10.7 (153)	123	3.3 (154)	113	11.5 (104)	125	4.0(114)	340	89.1 (56)	355	12.1 (166)
2555	8,109	50.2(139)	6,613	12.0 (146)	4,647	5.5 (137)	6,306	9.8 (118)	6,166	3.8(119)	10,472	81.1(139)	14,199	6.1 (75)
3001	1,086	48.4(145)	953	6.2 (164)	865	3.0 (160)	921	3.9 (155)	922	1.0(164)	2,491	78.6(156)	2,422	12.6 (171)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 89
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)		Number % (Rank)	
3002	280	49.3(142)	249 5.2 (165)	233 1.7 (165)	232 4.7 (151)	242 1.2(162)	724 89.0 (58)		655 5.5 (64)	
3003	6	* *	5 * *	5 * *	4 * *	5 * *	19 * *		25 * *	
3004	157	72.0 (22)	144 20.8 (65)	138 8.0 (100)	125 4.8 (150)	129 1.6(157)	569 72.6(168)		456 0.7 (2)	
3007	24	* *	30 * *	30 * *	27 * *	30 * *	114 86.8 (95)		95 * *	
3009	16	* *	9 * *	8 * *	8 * *	7 * *	33 * *		43 * *	
3010	2	* *	2 * *	3 * *	1 * *	1 * *	6 * *		7 * *	
3201	688	53.2(123)	617 14.6 (122)	509 6.3 (122)	605 14.7 (79)	597 4.9 (96)	1,303 94.2 (10)		1,481 7.2 (95)	
3301	4,654	64.8 (70)	3,896 19.5 (80)	3,211 9.7 (71)	3,499 9.3 (125)	3,637 4.3(105)	7,461 80.0(148)		9,608 5.8 (67)	
3302	326	60.4 (97)	318 13.8 (128)	289 8.0 (100)	277 7.9 (130)	304 3.6(122)	831 80.1(146)		864 5.4 (61)	
3304	171	95.3 (1)	129 60.5 (2)	127 40.9 (2)	108 36.1 (3)	111 18.9 (2)	407 81.1(139)		414 1.2 (7)	
3305	198	76.8 (13)	158 29.1 (13)	136 16.9 (8)	136 18.4 (46)	150 8.7 (45)	379 87.1 (85)		409 2.2 (12)	
3306	343	67.6 (47)	312 19.2 (86)	268 12.7 (26)	294 10.5 (109)	302 2.3(140)	635 93.5 (13)		726 5.8 (67)	
3308	84	* *	70 * *	68 * *	67 * *	66 * *	153 88.9 (60)		180 7.2 (95)	

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 90
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
3309	53	* *	44 * *	38 * *	41 * *	41 * *	149	85.2(111)	130	7.7 (108)
3550	96	* *	80 * *	69 * *	81 * *	76 * *	244	90.2 (39)	219	3.2 (20)
3551	28	* *	23 * *	24 * *	24 * *	23 * *	72	* *	70	* *
3552	54	* *	43 * *	46 * *	43 * *	42 * *	125	93.6 (12)	112	2.7 (15)
3701	1,339	63.7 (79)	1,169 16.7 (110)	1,054 9.1 (86)	1,063 19.4 (34)	1,116 11.0 (14)	2,630	94.3 (9)	2,954	7.7 (108)
3805	3,812	60.4 (97)	3,519 15.3 (119)	3,003 7.4 (109)	3,179 12.5 (95)	3,392 5.2 (93)	7,317	83.7(124)	8,358	7.6 (102)
3950	2,288	63.9 (76)	2,031 23.9 (35)	1,803 12.0 (35)	1,921 18.6 (45)	1,925 9.1 (34)	4,354	83.7(124)	5,035	5.4 (61)
3952	4	* *	11 * *	8 * *	8 * *	11 * *	42	* *	43	* *
3953	3	* *	4 * *	6 * *	2 * *	2 * *	50	* *	47	* *
3954	59	* *	57 * *	70 * *	51 * *	56 * *	264	88.6 (64)	223	0.9 (4)
4001	1,377	58.2(106)	1,094 15.3 (119)	971 6.1 (126)	1,037 12.3 (96)	987 5.3 (90)	2,828	63.9(184)	2,983	9.4 (141)
4002	1,281	63.2 (85)	1,098 19.9 (77)	967 10.1 (62)	1,069 14.8 (77)	1,036 5.7 (84)	2,605	69.6(175)	2,814	8.7 (132)
4103	114	53.5(122)	106 16.0 (116)	107 4.7 (143)	105 10.5 (109)	103 12.6 (9)	344	75.0(164)	319	6.6 (84)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 91
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
4106	2,130	61.3 (93)	1,773	19.3 (83)	1,435	9.3 (81)	1,678	8.6 (129)	1,652	4.8 (99)	3,919	79.9(149)	4,170	4.8 (40)
4107	3,764	65.4 (61)	3,401	22.6 (48)	2,703	10.5 (55)	3,108	16.5 (64)	3,194	9.2 (32)	6,890	87.0 (89)	8,033	4.1 (29)
4109	1,363	64.3 (74)	1,233	21.3 (57)	1,031	10.1 (62)	1,141	17.3 (56)	1,146	9.9 (25)	3,245	89.7 (47)	3,367	5.0 (50)
4110	1,088	63.5 (82)	1,228	20.0 (75)	1,126	9.7 (71)	1,021	5.6 (145)	1,087	3.4(124)	3,486	82.9(129)	3,379	5.6 (65)
4112	634	47.2(146)	562	27.0 (18)	569	14.9 (15)	529	3.8 (157)	549	2.0(150)	1,916	96.7 (3)	1,607	5.2 (54)
4113	733	67.1 (48)	626	22.7 (47)	552	12.5 (31)	593	5.9 (144)	546	2.6(134)	1,793	83.6(127)	1,707	1.4 (8)
4120	2,463	65.2 (67)	2,214	19.3 (83)	1,807	10.2 (58)	2,010	12.2 (98)	2,070	5.8 (80)	5,228	87.0 (89)	5,623	2.9 (16)
4160	2,246	66.7 (52)	2,001	26.3 (22)	1,611	14.6 (17)	1,781	15.7 (69)	1,803	11.1 (13)	4,209	84.9(116)	5,023	3.0 (17)
4351	661	68.1 (46)	594	20.2 (72)	557	11.3 (45)	558	19.2 (36)	567	9.0 (36)	1,549	88.4 (66)	1,555	10.7 (159)
4352	196	58.2(106)	169	17.2 (105)	161	9.3 (81)	154	11.0 (106)	163	4.3(105)	498	79.3(151)	481	18.5 (189)
4353	541	63.4 (83)	535	21.5 (56)	495	12.7 (26)	488	17.0 (59)	505	9.1 (34)	1,268	76.3(160)	1,264	12.6 (171)
4354	934	65.4 (61)	841	18.1 (98)	709	9.4 (79)	809	19.2 (36)	818	10.0 (23)	1,692	94.7 (6)	1,922	6.8 (89)
4355	379	66.5 (54)	331	20.2 (72)	304	7.6 (106)	298	19.1 (40)	308	4.2(109)	789	88.8 (61)	847	7.8 (112)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 92
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
4356	646	65.5 (58)	590	23.4 (40)	555	12.3 (34)	548	25.7 (13)	559	8.2 (47)	1,324	88.1 (71)	1,397	7.4 (98)
4357	1	* *	1	* *	0	* *	1	* *	1	* *	2	* *	3	* *
4358	203	79.3 (7)	183	35.0 (4)	177	24.3 (3)	169	11.8 (101)	177	2.8(131)	621	89.9 (42)	547	0.7 (2)
4359	621	74.7 (16)	491	23.2 (42)	501	10.2 (58)	447	14.8 (77)	470	5.7 (84)	1,536	81.4(135)	1,445	8.2 (119)
4360	207	77.3 (11)	176	19.3 (83)	158	7.0 (112)	166	18.7 (44)	169	5.9 (79)	435	89.9 (42)	455	11.6 (165)
4361	270	68.9 (40)	228	21.1 (61)	209	6.7 (117)	187	18.2 (47)	202	7.9 (53)	500	88.0 (72)	556	8.8 (133)
4362	369	77.0 (12)	309	26.9 (19)	310	13.5 (22)	259	23.9 (18)	274	12.0 (11)	775	89.7 (47)	826	12.2 (167)
4363	1,063	74.2 (17)	930	17.6 (103)	794	7.2 (111)	892	16.1 (66)	909	6.2 (74)	2,030	88.5 (65)	2,281	8.2 (119)
4401	1,599	65.5 (58)	1,400	17.1 (107)	1,152	10.1 (62)	1,313	14.9 (75)	1,338	6.0 (77)	2,699	92.4 (18)	3,263	6.4 (80)
4501	1,134	55.2(114)	1,001	14.2 (126)	901	8.2 (94)	922	13.2 (89)	933	3.8(119)	2,386	86.8 (95)	2,462	7.6 (102)
4502	1	* *	1	* *	0	* *	1	* *	1	* *	2	* *	1	* *
4591	0	* *	0	* *	0	* *	0	* *	0	* *	2	* *	2	* *
4601	1,946	67.1 (48)	1,646	20.8 (65)	1,321	9.4 (79)	1,510	14.0 (85)	1,549	5.0 (94)	3,044	79.6(150)	3,866	5.4 (61)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 93
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)	Smoking in Household
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
4701	1,693	68.3 (42)	1,303	22.0 (53)	1,213	8.8 (91)	1,205	18.9 (42)	1,258	7.5 (62)	2,631	90.7 (34)
5050	2,137	53.0(124)	1,687	14.3 (125)	1,472	5.8 (130)	1,606	10.2 (114)	1,548	4.6(101)	2,692	82.5(131)
5051	3,516	54.9(116)	2,701	15.9 (118)	2,434	7.0 (112)	2,679	10.5 (109)	2,540	5.0 (94)	4,526	80.1(146)
5052	3,283	51.0(134)	2,572	12.8 (142)	2,240	6.3 (122)	2,517	7.7 (131)	2,411	4.1(110)	4,134	78.9(153)
5101	2,880	65.3 (65)	2,376	20.0 (75)	2,031	10.2 (58)	2,315	12.0 (99)	2,218	4.9 (96)	4,130	85.1(113)
5202	765	68.2 (44)	607	25.2 (30)	529	13.4 (23)	598	19.2 (36)	588	9.0 (36)	1,215	90.5 (38)
5203	179	63.1 (87)	166	28.3 (14)	159	12.6 (29)	142	21.1 (25)	160	8.1 (48)	416	86.1(105)
5204	146	73.3 (19)	123	25.2 (30)	122	13.9 (20)	113	19.5 (33)	118	7.6 (59)	268	91.0 (31)
5601	1,455	71.3 (25)	1,308	22.8 (45)	1,116	9.3 (81)	1,238	24.0 (17)	1,256	9.0 (36)	2,739	89.4 (51)
5801	2,370	59.6(101)	2,127	14.9 (121)	1,803	6.9 (115)	1,970	12.7 (93)	2,049	5.8 (80)	3,911	82.6(130)
5803	266	69.5 (36)	235	23.8 (37)	192	12.0 (35)	231	20.8 (28)	228	8.3 (46)	433	85.0(114)
5804	101	50.5(136)	100	17.0 (108)	89	* *	99	* *	98	* *	232	86.6 (98)
6101	89	* *	80	* *	82	* *	79	* *	74	* *	233	90.1 (41)
											230	13.0 (174)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 94
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)										
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)		Number % (Rank)	
6102	237	66.7 (52)	207 25.1 (32)	220 12.7 (26)	204 22.5 (22)	196 7.7 (55)	678 89.2 (55)		634 6.9 (90)	
6103	4,248	52.0(129)	3,456 14.0 (127)	3,274 6.2 (125)	3,375 12.6 (94)	3,305 4.1(110)	8,547 84.8(117)		9,392 7.4 (98)	
6104	554	44.8(156)	337 13.1 (135)	268 4.9 (142)	340 9.4 (123)	324 2.5(135)	608 77.1(159)		897 13.3 (177)	
6105	76	* *	67 * *	74 * *	63 * *	64 * *	233 90.6 (36)		206 10.7 (159)	
6106	21	* *	17 * *	15 * *	19 * *	16 * *	38 * *		52 * *	
6301	5,845	54.1(117)	4,826 17.8 (101)	3,670 8.0 (100)	4,846 11.7 (103)	4,657 5.5 (87)	8,670 82.2(133)		11,436 4.1 (29)	
6312	3,584	63.3 (84)	2,956 24.1 (34)	2,152 11.8 (38)	2,936 14.9 (75)	2,848 7.5 (62)	4,906 79.2(152)		6,765 3.5 (24)	
6316	2,029	59.4(102)	1,716 17.8 (101)	1,284 9.2 (85)	1,655 13.2 (89)	1,620 5.3 (90)	2,628 81.3(137)		3,780 3.3 (23)	
6320	272	51.8(132)	253 22.5 (49)	201 13.9 (20)	233 12.0 (99)	243 6.6 (68)	663 88.8 (61)		695 1.4 (8)	
6325	182	62.6 (89)	180 17.2 (105)	140 5.7 (131)	166 15.7 (69)	174 6.3 (73)	358 86.0(106)		413 5.6 (65)	
6330	500	44.0(157)	421 11.9 (148)	314 2.9 (162)	416 6.7 (136)	410 2.7(133)	830 76.1(161)		1,054 8.3 (125)	
6335	671	50.1(140)	602 20.3 (71)	523 10.5 (55)	585 11.3 (105)	586 7.2 (65)	1,450 91.9 (22)		1,524 2.4 (13)	
6340	149	53.7(119)	130 20.8 (65)	106 16.0 (12)	120 10.0 (117)	125 8.0 (50)	380 94.7 (6)		388 1.0 (5)	

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 95
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)						Exclusive Breastfeeding (7)		≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
		Ever Breastfed		Breastfed At Least 6 Months		Breastfed At Least 12 Months		Breastfed At Least 3 Months		Breastfed At Least 6 Months		
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
6345	33	* *	36	* *	28	* *	33	* *	34	* *	82	* *
6350	333	45.3(154)	306	16.3 (113)	261	8.8 (91)	296	9.1 (127)	296	5.7 (84)	554	88.3 (70)
7304	239	52.7(126)	190	17.9 (99)	169	5.9 (129)	172	12.8 (92)	170	2.4(137)	591	87.0 (89)
7306	3,002	49.0(143)	2,318	12.9 (138)	2,001	3.6 (151)	2,084	8.9 (128)	2,021	4.5(102)	5,515	81.7(134)
7308	222	59.9(100)	170	18.2 (96)	144	7.6 (106)	153	11.8 (101)	129	5.4 (88)	457	89.1 (56)
7310	72	* *	48	* *	40	* *	46	* *	44	* *	85	* *
7401	3,026	56.6(112)	2,683	12.6 (143)	2,248	5.5 (137)	2,417	10.4 (113)	2,512	4.3(105)	5,159	86.7 (97)
7402	27	* *	41	* *	52	* *	28	* *	33	* *	133	91.0 (31)
7403	8	* *	15	* *	23	* *	9	* *	13	* *	53	* *
7404	7	* *	15	* *	14	* *	8	* *	13	* *	50	* *
7601	858	60.5 (94)	774	18.3 (95)	614	9.0 (89)	740	17.2 (58)	738	5.8 (80)	1,523	84.7(118)
7850	1,234	63.9 (76)	1,048	16.4 (111)	938	7.7 (105)	954	10.5 (109)	992	2.2(145)	2,477	90.7 (34)
7851	23	* *	17	* *	30	* *	11	* *	17	* *	143	96.5 (4)
											177	10.2 (150)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 96
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
7852	127	73.2 (20)	104	26.0 (25)	103	9.7 (71)	98	* *	103	4.9 (96)	236	92.4 (18)	276	7.6 (102)
8101	536	45.1(155)	176	13.1 (135)	85	* *	213	9.4 (123)	167	3.0(128)	226	69.9(173)	643	5.8 (67)
8102	51	* *	26	* *	26	* *	30	* *	26	* *	83	* *	90	* *
8105	138	59.4(102)	49	* *	27	* *	59	* *	46	* *	74	* *	166	4.2 (33)
8106	81	* *	31	* *	22	* *	30	* *	27	* *	89	* *	112	9.8 (144)
8150	51	* *	54	* *	53	* *	47	* *	49	* *	211	91.5 (25)	192	5.2 (54)
8151	3,597	68.3 (42)	3,075	28.0 (15)	2,482	15.1 (14)	2,923	17.3 (56)	2,877	10.3 (21)	5,596	80.9(141)	7,232	8.0 (116)
8152	474	60.5 (94)	370	11.1 (152)	286	3.1 (158)	352	7.7 (131)	349	2.3(140)	423	87.2 (84)	793	13.6 (178)
8153	31	* *	25	* *	24	* *	25	* *	22	* *	93	* *	87	* *
8154	381	79.5 (6)	303	64.7 (1)	263	41.4 (1)	274	48.2 (1)	254	39.4 (1)	726	96.8 (2)	857	1.4 (8)
8155	74	* *	62	* *	64	* *	46	* *	51	* *	246	87.0 (89)	210	10.5 (157)
8156	22	* *	22	* *	20	* *	15	* *	18	* *	83	* *	82	* *
8157	17	* *	16	* *	16	* *	16	* *	16	* *	78	* *	73	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 97
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
8201	3,373	56.6(112)	2,609	16.0 (116)	2,104	7.3 (110)	2,501	10.2 (114)	2,519	3.9(117)	2,510	85.4(109)	5,510	4.7 (38)
8202	3,653	62.9 (88)	2,815	27.5 (17)	2,421	11.0 (48)	2,687	6.1 (142)	2,642	3.7(121)	6,799	80.3(144)	7,474	4.9 (46)
8203	531	54.0(118)	417	16.3 (113)	343	7.6 (106)	392	10.7 (107)	386	3.9(117)	508	83.9(123)	873	6.1 (75)
8250	1,380	45.4(153)	933	12.4 (144)	666	8.1 (97)	954	6.5 (138)	856	4.1(110)	1,010	87.3 (83)	1,976	5.2 (54)
8252	73	* *	37	* *	29	* *	38	* *	31	* *	66	* *	106	4.7 (38)
8255	3,825	50.3(137)	2,796	14.5 (123)	2,439	6.6 (119)	2,673	9.7 (119)	2,729	4.1(110)	3,185	81.3(137)	6,385	6.3 (79)
8257	580	46.4(149)	473	25.6 (28)	413	12.6 (29)	456	4.6 (152)	453	2.9(130)	1,192	89.4 (51)	1,253	4.1 (29)
8258	1,664	47.0(147)	1,448	10.7 (153)	1,297	5.7 (131)	1,346	7.1 (134)	1,368	3.3(126)	2,435	85.7(108)	3,451	6.6 (84)
8260	828	71.5 (24)	654	31.2 (10)	577	20.8 (5)	620	6.1 (142)	629	2.5(135)	1,356	69.4(176)	1,592	5.3 (58)
8901	1,322	28.3(170)	1,079	8.0 (161)	844	3.3 (154)	1,084	3.8 (157)	1,013	1.8(153)	2,227	66.7(180)	2,726	13.2 (176)
8902	966	38.0(163)	752	12.2 (145)	560	5.7 (131)	759	6.2 (141)	706	2.3(140)	1,427	68.3(177)	1,850	10.9 (162)
8903	1,569	24.9(172)	1,266	7.3 (162)	1,033	2.7 (163)	1,274	1.9 (166)	1,195	1.3(161)	2,663	66.5(182)	3,202	12.2 (167)
8904	1,415	40.8(161)	1,113	13.6 (131)	823	6.9 (115)	1,108	6.5 (138)	1,034	2.1(147)	2,174	72.4(169)	2,740	10.7 (159)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 98
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
8905	3,488	36.2(167)	2,977	12.9 (138)	2,475	5.4 (140)	2,805	2.9 (161)	2,722	1.6(157)	7,049	75.9(162)	7,459	7.0 (93)
9004	8	* *	4	* *	3	* *	4	* *	4	* *	0	* *	9	* *
9010	6	* *	0	* *	0	* *	0	* *	0	* *	0	* *	6	* *
9012	51	* *	43	* *	40	* *	43	* *	42	* *	108	80.6(143)	93	* *
9013	134	45.5(152)	93	* *	73	* *	101	5.0 (148)	91	* *	148	65.5(183)	213	16.0 (183)
9015	41	* *	29	* *	23	* *	28	* *	29	* *	71	* *	75	* *
9016	189	50.8(135)	134	4.5 (168)	94	* *	150	2.7 (165)	134	0.7(165)	239	62.8(186)	330	16.1 (185)
9017	71	* *	46	* *	39	* *	41	* *	43	* *	114	78.1(158)	127	10.2 (150)
9018	151	50.3(137)	113	13.3 (134)	77	* *	119	6.7 (136)	111	1.8(153)	176	60.2(189)	243	8.2 (119)
9019	45	* *	35	* *	30	* *	31	* *	33	* *	86	* *	86	* *
9021	112	42.0(160)	76	* *	62	* *	71	* *	69	* *	141	80.9(141)	187	7.5 (100)
9040	4	* *	3	* *	2	* *	3	* *	3	* *	14	* *	20	* *
9041	130	42.3(159)	196	5.1 (166)	206	3.4 (152)	145	2.8 (163)	180	0.6(166)	716	74.0(165)	735	10.2 (150)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 99
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%	Number	%
		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
9045	1,789	46.6(148)	1,414	11.9 (148)	1,228	4.5 (145)	1,357	2.8 (163)	1,293	1.6(157)	3,541	69.8(174)	3,748	7.6 (102)
9050	1,971	34.2(168)	1,798	13.5 (132)	1,541	5.7 (131)	1,672	2.9 (161)	1,705	1.9(152)	3,971	66.7(180)	4,353	8.2 (119)
9051	1,931	52.5(127)	1,641	19.5 (80)	1,396	6.7 (117)	1,461	10.1 (116)	1,519	5.3 (90)	4,368	78.8(155)	4,433	3.0 (17)
9053	2,759	43.0(158)	2,316	11.3 (151)	1,676	5.5 (137)	2,260	4.5 (153)	2,264	2.3(140)	3,828	70.6(172)	5,444	17.5 (188)
9055	156	25.6(171)	87	* *	49	* *	96	* *	87	* *	1	* *	152	9.9 (145)
9058	845	64.7 (72)	740	16.1 (115)	704	8.2 (94)	715	3.4 (159)	724	1.7(156)	2,036	78.9(153)	1,895	9.1 (137)
9066	2,285	32.4(169)	2,074	5.1 (166)	1,676	1.8 (164)	2,022	3.2 (160)	2,045	1.2(162)	3,653	72.7(167)	4,701	8.4 (126)
9068	1,639	37.5(164)	1,420	8.5 (159)	1,151	3.9 (149)	1,382	3.9 (155)	1,374	2.3(140)	2,621	70.7(171)	3,303	13.0 (174)
9069	2,633	53.6(120)	2,240	9.7 (155)	1,579	4.2 (147)	2,219	5.0 (148)	2,203	2.0(150)	2,962	71.6(170)	4,752	10.2 (150)
9070	2,065	36.8(165)	1,697	9.1 (158)	1,420	3.3 (154)	1,573	7.6 (133)	1,579	2.8(131)	3,299	61.6(188)	4,168	8.6 (131)
9072	1,818	38.7(162)	1,516	8.4 (160)	1,217	3.0 (160)	1,495	5.6 (145)	1,482	2.2(145)	2,593	63.7(185)	3,541	11.2 (163)
9074	1,432	52.1(128)	1,301	12.0 (146)	974	4.2 (147)	1,274	5.3 (147)	1,275	1.8(153)	2,344	68.3(177)	3,076	10.1 (148)
9080	3,046	52.0(129)	2,502	18.5 (92)	2,219	8.2 (94)	2,481	4.5 (153)	2,395	2.1(147)	5,740	67.5(179)	6,488	7.8 (112)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pediatric Nutrition Surveillance (1)
Michigan

Page: 100
Run Date: 3 / 12 / 12

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)									
	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)	Smoking in Household	
Clinic	Number	% Ever Breastfed	Number % Breastfed At Least 6 Months (Rank)	Number % Breastfed At Least 12 Months (Rank)	Number % Breastfed At Least 3 Months (Rank)	Number % Breastfed At Least 6 Months (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)
9151	0	* *	0 * *	1 * *	0 * *	0 * *	1 * *	2 * *	
Unknown Clinics	298	55.0(115)	187 26.2 (24)	151 6.0 (127)	202 17.8 (52)	178 13.5 (7)	412 62.6(187)	503 20.3 (193)	
Michigan	200,166	57.3	168,817 17.7	142,897 8.5	158,974 12.2	159,216 5.5	353,084 82.4	418,132 7.0	

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

